

# Methylxanthines and Nicotine

Chip Gresham  
Daniel E. Brooks

## REFERENCES

1. Sankar J, Lodha R, Kabra SK: Doxofylline: the next generation methylxanthine. *Indian J Pediatr* 75: 251, 2008. [PMID: 18376093]
2. Shan D, Wu HM, Yuan QY, et al: Pentoxyfylline for diabetic kidney disease. *Cochrane Database Syst Rev* 2: CD006800, 2012. [PMID: 22336824]
3. Temple JL: Caffeine use in children: what we know, what we have left to learn, and why we should worry. *Neurosci Biobehav Rev* 33: 793, 2009. [PMID: 19428492]
4. Wolk BJ, Ganetsky M, Babu KM: Toxicity of energy drinks. *Curr Opin Pediatr* 58: 243, 2012. [PMID: 22426157]
5. Babu KM, Church RJ, Lewander W: Energy drinks: the new eye-opener for adolescents. *Clin Pediatr Emerg Med* 9: 35, 2008. [No PMID]
6. Kupper TE, Strohl KP, Hoefer M, et al: Low-dose theophylline reduces symptoms of acute mountain sickness. *J Travel Med* 15: 307, 2008. [PMID: 19006503]
7. Dai B, Liu Y, Fu L, et al: Effect of theophylline on prevention of contrast-induced acute kidney injury: a meta-analysis of randomized controlled trials. *Am J Kidney Dis* 60: 360, 2012. [PMID: 22516682]
8. Patanè S, Marte F, La Rosa FC, Rocca RL: Atrial fibrillation associated with chocolate intake abuse and chronic salbutamol inhalation abuse. *Int J Cardiol* 145: e74, 2010. [PMID: 19171401]
9. Minton NA, Henry JA: Acute and chronic human toxicity of theophylline. *Hum Exp Toxicol* 15: 471, 1996. [PMID: 8793529]
10. Babu KM, Zuckerman MD, Cherkes JK, Hack JB: First-onset seizure after use of an energy drink [corrected]. *Pediatr Emerg Care* 27: 539, 2011. [Erratum in *Pediatr Emerg Care* 28: 210, 2012] [PMID: 21642791]
11. Di Rocco JR, During A, Morelli PJ, et al: Atrial fibrillation in healthy adolescents after highly caffeinated beverage consumption: two case reports. *J Med Case Rep* 5: 18, 2011. [PMID: 21247417]
12. Nagajothi N, Khraisat A, Velazquez-Cecena JL, et al: Energy drink-related supraventricular tachycardia. *Am J Med* 121: e4, 2008. [PMID: 18374674]
13. Sessler CN, Cohen MD: Cardiac arrhythmias during theophylline toxicity. A prospective continuous electrocardiographic study. *Chest* 98: 672, 1990. [PMID: 2394145]
14. Mitchell ES, Slettenaar M, vd Meer N, et al: Differential contributions of theobromine and caffeine on mood, psychomotor performance and blood pressure. *Physiol Behav* 104: 816, 2011. [PMID: 21839757]
15. Rudolph T, Knudsen K: A case of fatal caffeine poisoning. *Acta Anaesthesiol Scand* 54: 521, 2010. [PMID: 20096021]
16. Shannon M: Life-threatening events after theophylline overdose: a 10-year prospective analysis. *Arch Intern Med* 159: 989, 1999. [PMID: 10326941]
17. Chyka PA, Seger D, Krenzelok EP, et al: Position paper: single-dose activated charcoal. *Clin Toxicol (Phila)* 43: 61, 2005. [PMID: 15822758]
18. [No authors listed]: Position statement and practice guidelines on the use of multi-dose activated charcoal in the treatment of acute poisoning. American Academy of Clinical Toxicology; European Association of Poisons Centres and Clinical Toxicologists. *J Toxicol Clin Toxicol* 37: 731, 1999. [PMID: 10584586]
19. [No authors listed]: Position paper: whole bowel irrigation. *J Toxicol Clin Toxicol* 42: 843, 2004. [PMID: 15533024]
20. Krenzelok EP, McGuigan M, Lheur P: Position statement: ipecac syrup. American Academy of Clinical Toxicology; European Association of Poisons Centres and Clinical Toxicologists. *J Toxicol Clin Toxicol* 35: 699, 1997. [PMID: 9482425]
21. Krenzelok E, Vale A: Position statements: gut decontamination. American Academy of Clinical Toxicology; European Association of Poisons Centres and Clinical Toxicologists. *J Toxicol Clin Toxicol* 35: 695, 1997. [PMID: 9482424]
22. Gallop K: Review article: phenytoin use and efficacy in the ED. *Emerg Med Australas* 22: 108, 2010. [PMID: 20534046]
23. Huerta C, Lanes SF, García Rodríguez LA: Respiratory medications and the risk of cardiac arrhythmias. *Epidemiology* 16: 360, 2005. [PMID: 15824553]
24. Kneser J, Wehmeier P, Lichtenhagen R, Hooper MM, Kielstein JT: Successful treatment of life-threatening theophylline intoxication in a pregnant patient by hemodialysis. *Clin Nephrol* [Epub ahead of print] 2012 Jul 12. [PMID: 22541679]
25. Kapur R, Smith MD: Treatment of cardiovascular collapse from caffeine overdose with lidocaine, phenylephrine, and hemodialysis. *Am J Emerg Med* 2: 253, 2009. [PMID: 19371553]
26. Eden G, Busch M, Kühn-Velten WN, Schneider A, Kielstein JT: Successful treatment of life-threatening pentoxyfylline intoxication by high-flux hemodialysis. *Clin Nephrol* 75: 171, 2011. [PMID: 2125548]
27. Shannon M: Predictors of major toxicity after theophylline overdose. *Ann Intern Med* 119: 1161, 1993. [PMID: 8239246]
28. The National Institute on Drug Abuse: *Research Report Series—Tobacco Addiction*. NIH Publication No. 06-4342. Bethesda, MD: National Institutes of Health, July 2012.
29. Appleton S: Frequency and outcomes of accidental ingestion of tobacco products in young children. *Regul Toxicol Pharmacol* 61: 210, 2011. [PMID: 21821089]
30. Arcury TA, Quandt SA, Preisser JS, Bernert JT, Norton D, Wang J: High levels of transdermal nicotine exposure produce green tobacco sickness in Latino farmworkers. *Nicotine Tob Res* 5: 315, 2003. [PMID: 12791526]
31. Shiffman S, Fant RV, Buchhalter AR, et al: Nicotine delivery systems. *Expert Opin Drug Deliv* 2: 563, 2005. [PMID: 16296775]
32. Smolinske SC, Spoerke DG, Spiller SK, et al: Cigarette and nicotine chewing gum toxicity in children. *Hum Toxicol* 7: 27, 1988. [PMID: 3346035]
33. Wong S, Ordean A, Kahan M; Maternal Fetal Medicine Committee; Family Physicians Advisory Committee; Medico-Legal Committee; Society of Obstetricians and Gynaecologists of Canada: Substance use in pregnancy. *J Obstet Gynaecol Can* 33: 367, 2011. [PMID: 21501542]
34. Morrell HE, Cohen LM, al'Absi M: Physiological and psychological symptoms and predictors in early nicotine withdrawal. *Pharmacol Biochem Behav* 89: 272, 2008. [PMID: 18294681]