

## 34 Hemodynamic Monitoring



### REFERENCES

1. Wo CC, Shoemaker WC, Appel PL, et al: Unreliability of blood pressure and heart rate to evaluate cardiac output in emergency resuscitation and critical illness. *Crit Care Med* 21: 218, 1993.
2. Swan HJ, Ganz W, Forrester J, et al: Catheterization of the heart in man with use of a flow-directed balloon-tipped catheter. *N Engl J Med* 283: 447, 1970.
3. Shure D: Pulmonary-artery catheters—peace at last? *N Engl J Med* 354: 2273, 2006.
4. Eisenberg PR, Jaffe AS, Schuster DP: Clinical evaluation compared to pulmonary artery catheterization in the hemodynamic assessment of critically ill patients. *Crit Care Med* 12: 549, 1984.
5. Connors AF Jr, McCaffree DR, Gray BA: Evaluation of right-heart catheterization in the critically ill patient without acute myocardial infarction. *N Engl J Med* 308: 263, 1983.
6. Hall WD: Stephen Hales: theologian, botanist, physiologist, discoverer of hemodynamics. *Clin Cardiol* 10: 487, 1987.
7. Pickering TG, Hall JE, Appel LJ, et al: Recommendations for blood pressure measurement in humans and experimental animals: part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Circulation* 111: 697, 2005.
8. Deakin CD, Low JL: Accuracy of the advanced trauma life support guidelines for predicting systolic blood pressure using carotid, femoral, and radial pulses: observational study. *BMJ* 321: 673, 2000.
9. Brown WC, O'Brien ET, Semple PF: The sphygmomanometer of Riva-Rocci 1896–1996. *J Hum Hypertens* 10: 723, 1996.
10. Segall HN: How Korotkoff, the surgeon, discovered the auscultatory method of measuring arterial pressure. *Ann Intern Med* 83: 561, 1975.
11. Karnath B: Sources of error in blood pressure measurement. *Hosp Physician* 38: 33, 2002.
12. Marey EJ: *Pression et vitesse du sang. Pratique des hautes études de M Marey. Physiologie Experimentale* Paris, 1876.
13. Pickering TG: Principles and techniques of blood pressure measurement. *Cardiol Clin* 20: 207, 2002.
14. Pickering TG: Blood pressure variability and ambulatory monitoring. *Curr Opin Nephrol Hypertens* 2: 380, 1993.
15. Cohn JN: Blood pressure measurement in shock. Mechanism of inaccuracy in auscultatory and palpatory methods. *JAMA* 199: 118, 1967.
16. Antman EM, Anbe DT, Armstrong PW, et al: ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Revise the 1999 Guidelines for the Management of Patients with Acute Myocardial Infarction). *Circulation* 110: e82, 2004.
17. Antonelli M, Levy M, Andrews PJ, et al: Hemodynamic monitoring in shock and implications for management. International Consensus Conference, Paris, France, 27–28 April 2006. *Intensive Care Med* 33: 575, 2007.
18. Scheer B, Perel A, Pfeiffer UJ: Clinical review: complications and risk factors of peripheral arterial catheters used for haemodynamic monitoring in anaesthesia and intensive care medicine. *Crit Care* 6: 199, 2002.
19. Mohr R, Lavee J, Goor DA: Inaccuracy of radial artery pressure measurement after cardiac operations. *J Thorac Cardiovasc Surg* 94: 286, 1987.
20. Barone JE, Madlinger RV: Should an Allen test be performed before radial artery cannulation? *J Trauma* 61: 468, 2006.
21. LeDoux D, Astiz ME, Carpati CM, Rackow EC: Effects of perfusion pressure on tissue perfusion in septic shock. *Crit Care Med* 28: 2729, 2000.
22. Bourgoin A, Leone M, Delmas A, et al: Increasing mean arterial pressure in patients with septic shock: effects on oxygen variables and renal function. *Crit Care Med* 33: 780, 2005.
23. Bratton SL, Chestnut RM, Ghajar J, et al: Guidelines for the management of severe traumatic brain injury. IX. Cerebral perfusion thresholds. *J Neurotrauma* 24(Suppl 1): S59, 2007.
24. Bratton SL, Chestnut RM, Ghajar J, et al: Guidelines for the management of severe traumatic brain injury. I. Blood pressure and oxygenation. *J Neurotrauma* 24(Suppl 1): S7, 2007.
25. Struchen MA, Hannay HJ, Contant CF, Robertson CS: The relation between acute physiological variables and outcome on the Glasgow Outcome Scale and Disability Rating Scale following severe traumatic brain injury. *J Neurotrauma* 18: 115, 2001.
26. Stern SA, Dronen SC, Birrer P, Wang X: Effect of blood pressure on hemorrhage volume and survival in a near-fatal hemorrhage model incorporating a vascular injury. *Ann Emerg Med* 22: 155, 1993.
27. Bickell WH, Wall MJ Jr, Pepe PE, et al: Immediate versus delayed fluid resuscitation for hypotensive patients with penetrating torso injuries. *N Engl J Med* 331: 1105, 1994.
28. Landis EM, Brown E, Fauteux M, Wise C: Central venous pressure in relation to cardiac “competence” blood volume and exercise. *J Clin Invest* 25: 237, 1946.
29. Marik PE, Baram M, Vahid B: Does central venous pressure predict fluid responsiveness? A systematic review of the literature and the tale of seven maids. *Chest* 134: 172, 2008.
30. McGee DC, Gould MK: Preventing complications of central venous catheterization. *N Engl J Med* 348: 1123, 2003.
31. Lorente L, Henry C, Martin MM, et al: Central venous catheter-related infection in a prospective and observational study of 2,595 catheters. *Crit Care* 9: R631, 2005.
32. Miller AH, Roth BA, Mills TJ, et al: Ultrasound guidance versus the landmark technique for the placement of central venous catheters in the emergency department. *Acad Emerg Med* 9: 800, 2002.
33. Pronovost P, Needham D, Berenholtz S, et al: An intervention to decrease catheter-related bloodstream infections in the ICU. *N Engl J Med* 355: 2725, 2006.
34. Parienti JJ, Thirion M, Megarbane B, et al: Femoral vs jugular venous catheterization and risk of nosocomial events in adults requiring acute renal replacement therapy: a randomized controlled trial. *JAMA* 299: 2413, 2008.
35. Magder S: Central venous pressure: a useful but not so simple measurement. *Crit Care Med* 34: 2224, 2006.
36. Desmond J, Megahed M: Is the central venous pressure reading equally reliable if the central line is inserted via the femoral vein. *Emerg Med J* 20: 467, 2003.
37. Leonard AD, Allsager CM, Parker JL, et al: Comparison of central venous and external jugular venous pressures during repair of proximal femoral fracture. *Br J Anaesth* 101: 166, 2008.
38. Charalambous C, Barker TA, Zipitis CS, et al: Comparison of peripheral and central venous pressures in critically ill patients. *Anaesth Intensive Care* 31: 34, 2003.
39. Constant J: Using internal jugular pulsations as a manometer for right atrial pressure measurements. *Cardiology* 93: 26, 2000.
40. Drazner MH, Rame JE, Stevenson LW, Dries DL: Prognostic importance of elevated jugular venous pressure and a third heart sound in patients with heart failure. *N Engl J Med* 345: 574, 2001.
41. Lipton B: Estimation of central venous pressure by ultrasound of the internal jugular vein. *Am J Emerg Med* 18: 432, 2000.
42. Randazzo MR, Snoey ER, Levitt MA, Binder K: Accuracy of emergency physician assessment of left ventricular ejection fraction and central venous pressure using echocardiography. *Acad Emerg Med* 10: 973, 2003.
43. Ward KR, Tiba MH, Barbee RW, et al: A new noninvasive method to determine central venous pressure. *Resuscitation* 70: 238, 2006.
44. Gelman S: Venous function and central venous pressure: a physiologic story. *Anesthesiology* 108: 735, 2008.
45. Oohashi S, Endoh H: Does central venous pressure or pulmonary capillary wedge pressure reflect the status of circulating blood volume in patients after extended transthoracic esophagectomy? *J Anesth* 19: 21, 2005.
46. Kumar A, Anel R, Bunnell E, et al: Pulmonary artery occlusion pressure and central venous pressure fail to predict ventricular filling volume, cardiac performance, or the response to volume infusion in normal subjects. *Crit Care Med* 32: 691, 2004.
47. Weil MH, Shubin H, Rosoff L: Fluid repletion in circulatory shock: central venous pressure and other practical guides. *JAMA* 192: 668, 1965.
48. Rivers E, Nguyen B, Havstad S, et al: Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 345: 1368, 2001.
49. Hollenberg SM, Ahrens TS, Annane D, et al: Practice parameters for hemodynamic support of sepsis in adult patients: 2004 update. *Crit Care Med* 32: 1928, 2004.
50. Westphal GA, Silva E, Caldeira Filho M, et al: Variation in amplitude of central venous pressure curve induced by respiration is a useful tool to reveal fluid responsiveness in postcardiac surgery patients. *Shock* 26: 140, 2006.
51. Magder SA, Georgiadis G, Cheong T: Respiratory variations in right atrial pressure predict response to fluid challenge. *J Crit Care* 7: 76, 1992.
52. Kong R, Singer M: Insertion of a pulmonary artery flotation catheter: how to do it. *Br J Hosp Med* 57: 432, 1997.
53. Mihm FG, Gettinger A, Hanson CW 3rd, et al: A multicenter evaluation of a new continuous cardiac output pulmonary artery catheter system. *Crit Care Med* 26: 1346, 1998.
54. Connors AF Jr, Sproff T, Dawson NV, et al: The effectiveness of right heart catheterization in the initial care of critically ill patients. SUPPORT Investigators. *JAMA* 276: 889, 1996.
55. Shah MR, Hasselblad V, Stevenson LW, et al: Impact of the pulmonary artery catheter in critically ill patients: meta-analysis of randomized clinical trials. *JAMA* 294: 1664, 2005.
56. Wheeler AP, Bernard GR, Thompson BT, et al: Pulmonary-artery versus central venous catheter to guide treatment of acute lung injury. *N Engl J Med* 354: 2213, 2006.
57. Johnston IG, Jane R, Fraser JE, et al: Survey of intensive care nurses' knowledge relating to the pulmonary artery catheter. *Anaesth Intensive Care* 32: 564, 2004.
58. Trottier SJ, Taylor RW: Physicians' attitudes toward and knowledge of the pulmonary artery catheter: Society of Critical Care Medicine membership survey. *New Horiz* 5: 201, 1997.
59. Vincent JL, Pinsky MR, Sprung CL, et al: The pulmonary artery catheter: in medio virtus. *Crit Care Med* 36: 3093, 2008.
60. Wertheimer L: Non-invasive techniques in the diagnosis and management of acute cardiac problems. *Med Clin North Am* 57: 1491, 1973.
61. Velmahos GC, Wo CC, Demetriades D, Shoemaker WC: Early continuous noninvasive hemodynamic monitoring after severe blunt trauma. *Injury* 30: 209, 1999.
62. Shoemaker WC, Belzberg H, Wo CC, et al: Multicenter study of noninvasive monitoring systems as alternatives to invasive monitoring of acutely ill emergency patients. *Chest* 114: 1643, 1988.
63. Marik PE, Baram M: Noninvasive hemodynamic monitoring in the intensive care unit. *Crit Care Clin* 23: 383, 2007.
64. Summers RL, Shoemaker WC, Peacock WF, et al: Bench to bedside: electrophysiologic and clinical principles of noninvasive hemodynamic monitoring using impedance cardiography. *Acad Emerg Med* 10: 669, 2003.
65. Newman DG, Callister R: The non-invasive assessment of stroke volume and cardiac output by impedance cardiography: a review. *Aviat Space Environ Med* 70: 780, 1999.
66. Raaijmakers E, Faes TJ, Scholten RJ, et al: A meta-analysis of three decades of validating thoracic impedance cardiography. *Crit Care Med* 27: 1203, 1999.
67. Peacock WF, Summers RL, Vogel J, Emerman CE: Impact of impedance cardiography on diagnosis and therapy of emergent dyspnea: the ED-IMPACT trial. *Acad Emerg Med* 13: 365, 2006.

68. Lo HY, Liao SC, Ng CJ, et al: Utility of impedance cardiography for dyspneic patients in the ED. *Am J Emerg Med* 25: 437, 2007.
69. Keren H, Burkhoff D, Squara P: Evaluation of a noninvasive continuous cardiac output monitoring system based on thoracic bioimpedance. *Am J Physiol Heart Circ Physiol* 293: H583, 2007.
70. Raval NY, Squara P, Cleman M, et al: Multicenter evaluation of noninvasive cardiac output measurement by bioimpedance technique. *J Clin Monit Comput* 22: 113, 2008.
71. Huntsman LL, Stewart DK, Barnes SR, et al: Noninvasive Doppler determination of cardiac output in man. Clinical validation. *Circulation* 67: 593, 1983.
72. Dark PM, Singer M: The validity of trans-esophageal Doppler ultrasonography as a measure of cardiac output in critically ill adults. *Intensive Care Med* 30: 2060, 2004.
73. Mythen MG, Webb AR: Perioperative plasma volume expansion reduces the incidence of gut mucosal hypoperfusion during cardiac surgery. *Arch Surg* 130: 423, 1995.
74. Sinclair S, James S, Singer M: Intraoperative intravascular volume optimisation and length of hospital stay after repair of proximal femoral fracture: randomised controlled trial. *BMJ* 315: 909, 1997.
75. Gan TJ, Soppitt A, Maroof M, et al: Goal-directed intraoperative fluid administration reduces length of hospital stay after major surgery. *Anesthesiology* 97: 820, 2002.
76. McKendry M, McGloin H, Saberi D, et al: Randomised controlled trial assessing the impact of a nurse delivered, flow monitored protocol for optimisation of circulatory status after cardiac surgery. *BMJ* 329: 258, 2004.
77. Noblett SE, Snowden CP, Shenton BK, Horgan AF: Randomized clinical trial assessing the effect of Doppler-optimized fluid management on outcome after elective colorectal resection. *Br J Surg* 93: 1069, 2006.
78. Rodriguez RM, Lum-Lung M, Dixon K, Nothmann A: A prospective study on esophageal Doppler hemodynamic assessment in the ED. *Am J Emerg Med* 24: 658, 2006.
79. Venn R, Steele A, Richardson P, et al: Randomized controlled trial to investigate influence of the fluid challenge on duration of hospital stay and perioperative morbidity in patients with hip fractures. *Br J Anaesth* 88: 65, 2002.
80. van Lelyveld-Haas LE, van Zanten AR, Borm GF, Tjan DH: Clinical validation of the non-invasive cardiac output monitor USCOM-1A in critically ill patients. *Eur J Anaesthesiol* 25:917, 2008.
81. Wong LS, Yong BH, Young KK, et al: Comparison of the USCOM ultrasound cardiac output monitor with pulmonary artery catheter thermodilution in patients undergoing liver transplantation. *Liver Transpl* 14: 1038, 2008.
82. Nguyen HB, Losey T, Rasmussen J, et al: Interrater reliability of cardiac output measurements by transcutaneous Doppler ultrasound: implications for noninvasive hemodynamic monitoring in the ED. *Am J Emerg Med* 24: 828, 2006.
83. Stewart GM, Nguyen HB, Kim TY, et al: Inter-rater reliability for noninvasive measurement of cardiac function in children. *Pediatr Emerg Care* 24: 433, 2008.
84. Dey I, Sprivilis P: Emergency physicians can reliably assess emergency department patient cardiac output using the USCOM continuous wave Doppler cardiac output monitor. *Emerg Med Australas* 17: 193, 2005.
85. Knobloch K, Tepe J, Rossner D, et al: Combined NT-pro-BNP and CW-Doppler ultrasound cardiac output monitoring (USCOM) in epirubicin and liposomal doxorubicin therapy. *Int J Cardiol* 128: 316, 2008.
86. Knobloch K, Tepe J, Lichtinghagen R, et al: Simultaneous hemodynamic and serological cardiotoxicity monitoring during immunotherapy with trastuzumab. *Int J Cardiol* 125: 113, 2008.
87. Knobloch K, Hubrich V, Rohmann P, et al: Non-invasive determination of cardiac output by continuous wave Doppler in air rescue service. *Anesthesiol Intensivmed Notfallmed Schmerzther* 40: 750, 2005.
88. Linton RAF, Band DM, Haire KM: A new method of measuring cardiac output in man using lithium dilution. *Br J Anaesth* 71: 262, 1993.
89. Jonas MM, Kelly FE, Linton RA, et al: A comparison of lithium dilution cardiac output measurements made using central and antecubital venous injection of lithium chloride. *J Clin Monit Comput* 15: 525, 1999.
90. Della Rocca G, Costa MG, Pompei L, et al: Continuous and intermittent cardiac output measurement: pulmonary artery catheter versus aortic transpulmonary technique. *Br J Anaesth* 88: 350, 2002.
91. Opdam HI, Wan L, Bellomo R: A pilot assessment of the FloTrac cardiac output monitoring system. *Intensive Care Med* 33: 344, 2007.
92. Lorsomradee S, Lorsomradee SR, Cromheecke S, De Hert SG: Continuous cardiac output measurement: arterial pressure analysis versus thermodilution technique during cardiac surgery with cardiopulmonary bypass. *Anaesthesia* 62: 979, 2007.
93. de Waal EE, Kalkman CJ, Rex S, Buhre WF: Validation of a new arterial pulse contour-based cardiac output device. *Crit Care Med* 35: 1904, 2007.
94. McGee WT, Horswell JL, Calderon J, et al: Validation of a continuous, arterial pressure-based cardiac output measurement: a multicenter, prospective clinical trial. *Crit Care* 11: R105, 2007.
95. Starling E: The Linacre Lecture on the law of the heart given at Cambridge, 1915. London, UK, Longmans, Green and Co., 1918.
96. Michard F, Teboul JL: Predicting fluid responsiveness in ICU patients: a critical analysis of the evidence. *Chest* 121: 2000, 2002.
97. Coudray A, Romand JA, Treggiari M, Bendjelid K: Fluid responsiveness in spontaneously breathing patients: a review of indexes used in intensive care. *Crit Care Med* 33: 2757, 2005.
98. Hadian M, Pinsky MR: Functional hemodynamic monitoring. *Curr Opin Crit Care* 13: 318, 2007.
99. Pinsky MR, Payen D: Functional hemodynamic monitoring. *Crit Care* 9: 566, 2005.
100. Michard F, Boussat S, Chemla D, et al: Relation between respiratory changes in arterial pulse pressure and fluid responsiveness in septic patients with acute circulatory failure. *Am J Respir Crit Care Med* 162: 134, 2000.
101. Lopes MR, Oliveira MA, Pereira VO, et al: Goal-directed fluid management based on pulse pressure variation monitoring during high-risk surgery: a pilot randomized controlled trial. *Crit Care* 11: R100, 2007.
102. Pinsky MR: Functional hemodynamic monitoring. *Intensive Care Med* 28: 386, 2002.
103. Pinsky MR, Vincent JL: Let us use the pulmonary artery catheter correctly and only when we need it. *Crit Care Med* 33: 1119, 2005.
104. Monnet X, Rienzo M, Osman D, et al: Passive leg raising predicts fluid responsiveness in the critically ill. *Crit Care Med* 34: 1402, 2006.
105. Lee J, Wright F, Barber R, Stanley L: Central venous oxygen saturation in shock: a study in man. *Anesthesiology* 36: 472, 1972.
106. Scheinman MM, Brown MA, Rapaport E: Critical assessment of use of central venous oxygen saturation as a mirror of mixed venous oxygen in severely ill cardiac patients. *Circulation* 40: 165, 1969.
107. Reinhart K, Kuhn HJ, Hartog C, Bredele DL: Continuous central venous and pulmonary artery oxygen saturation monitoring in the critically ill. *Intensive Care Med* 30: 1572, 2004.
108. Tahvanainen J, Meretoja O, Nikki P: Can central venous blood replace mixed venous blood samples? *Crit Care Med* 10: 758, 1982.
109. Dueck MH, Klimek M, Appenrodt S, et al: Trends but not individual values of central venous oxygen saturation agree with mixed venous oxygen saturation during varying hemodynamic conditions. *Anesthesiology* 103: 249, 2005.
110. Cortez A, Zito J, Lucas CE, Gerrick SJ: Mechanism of inappropriate polyuria in septic patients. *Arch Surg* 112: 471, 1977.
111. Rady MY, Rivers EP, Nowak RM: Resuscitation of the critically ill in the ED: responses of blood pressure, heart rate, shock index, central venous oxygen saturation, and lactate. *Am J Emerg Med* 14: 218, 1996.
112. Marshall JC: Inflammation, coagulopathy, and the pathogenesis of multiple organ dysfunction syndrome. *Crit Care Med* 29: S99, 2001.
113. Gattinoni L, Brazzi L, Pelosi P, et al: A trial of goal-oriented hemodynamic therapy in critically ill patients. SVO<sub>2</sub> Collaborative Group. *N Engl J Med* 333: 1025, 1995.
114. Hayes MA, Yau EH, Timmins AC, et al: Response of critically ill patients to treatment aimed at achieving supranormal oxygen delivery and consumption. Relationship to outcome. *Chest* 103: 886, 1993.
115. Trzeciak S, Dellinger RP, Abate NL, et al: Translating research to clinical practice: a 1-year experience with implementing early goal-directed therapy for septic shock in the emergency department. *Chest* 129: 225, 2006.
116. Shapiro NI, Howell MD, Talmor D, et al: Implementation and outcomes of the Multiple Urgent Sepsis Therapies (MUST) protocol. *Crit Care Med* 34: 1025, 2006.
117. Jones AE, Shapiro NI, Roshon M: Implementing early goal-directed therapy in the emergency setting: the challenges and experiences of translating research innovations into clinical reality in academic and community settings. *Acad Emerg Med* 14:1072, 2007.
118. Nguyen HB, Corbett SW, Steele R, et al: Implementation of a bundle of quality indicators for the early management of severe sepsis and septic shock is associated with decreased mortality. *Crit Care Med* 35: 1105, 2007.
119. Kompanje EJ, Jansen TC, van der Hoven B, Bakker J: The first demonstration of lactic acid in human blood in shock by Johann Joseph Scherer (1814–1869) in January 1843. *Intensive Care Med* 33: 1967, 2007.
120. Huckabee WE: Abnormal resting blood lactate. I. The significance of hyperlactemia in hospitalized patients. *Am J Med* 30: 833, 1961.
121. Broder G, Weil MH: Excess lactate: an index of reversibility of shock in human patients. *Science* 143: 1457, 1964.
122. Peretz DI, Scott HM, Duff J, et al: The significance of lactic acidemia in the shock syndrome. *Ann N Y Acad Sci* 119: 1133, 1965.
123. Levy B, Gibot S, Franck P, et al: Relation between muscle Na<sup>+</sup>K<sup>+</sup> ATPase activity and raised lactate concentrations in septic shock: a prospective study. *Lancet* 365: 871, 2005.
124. Gore DC, Jahoor F, Hibbert JM, et al: Lactic acidosis during sepsis is related to increased pyruvate production, not deficits in tissue oxygen availability. *Ann Surg* 224: 97, 1996.
125. Vary TC, Siegel JH, Nakatani T, et al: Effect of sepsis on activity of pyruvate dehydrogenase complex in skeletal muscle and liver. *Am J Physiol* 250: E634, 1986.
126. Fall PJ, Szerlip HM: Lactic acidosis: from sour milk to septic shock. *J Intensive Care Med* 20: 255, 2005.
127. Kruse JA, Zaidi SA, Carlson RW: Significance of blood lactate levels in critically ill patients with liver disease. *Am J Med* 83: 77, 1987.
128. Levraut J, Bouatirou T, Ichai C, et al: Reliability of anion gap as an indicator of blood lactate in critically ill patients. *Intensive Care Med* 23: 417, 1997.
129. Iberti TJ, Leibowitz AB, Papadakis PJ, Fischer EP: Low sensitivity of the anion gap as a screen to detect hyperlactatemia in critically ill patients. *Crit Care Med* 18: 275, 1990.
130. Weil MH, Michaels S, Rackow EC: Comparison of blood lactate concentrations in central venous, pulmonary artery, and arterial blood. *Crit Care Med* 15: 489.
131. Murdoch IA, Turner C, Dalton RN: Arterial or mixed venous lactate measurement in critically ill children. Is there a difference? *Acta Paediatr* 83: 412, 1994.
132. Gallagher EJ, Rodriguez K, Touger M: Agreement between peripheral venous and arterial lactate levels. *Ann Emerg Med* 29: 479, 1997.
133. Weil MH, Afifi AA: Experimental and clinical studies on lactate and pyruvate as indicators of the severity of acute circulatory failure (shock). *Circulation* 41: 989, 1970.
134. Bakker J, Coffernils M, Leon M, et al: Blood lactate levels are superior to oxygen-derived variables in predicting outcome in human septic shock. *Chest* 99: 956, 1991.
135. Aduen J, Bernstein WK, Khastgir T, et al: The use and clinical importance of a substrate-specific electrode for rapid determination of blood lactate concentrations. *JAMA* 272: 1678, 1994.

136. Shapiro NI, Howell MD, Talmor D, et al: Serum lactate as a predictor of mortality in emergency department patients with infection. *Ann Emerg Med* 45: 524, 2005.
137. Trzeciak S, Dellinger RP, Chansky ME, et al: Serum lactate as a predictor of mortality in patients with infection. *Intensive Care Med* 33: 970, 2007.
138. Vincent JL, Dufaye P, Berre J, et al: Serial lactate determinations during circulatory shock. *Crit Care Med* 11: 449, 1983.
139. Abramson D, Scalea TM, Hitchcock R, et al: Lactate clearance and survival following injury. *J Trauma* 35: 584, 1993.
140. Bakker J, Gris P, Coffernils M, et al: Serial blood lactate levels can predict the development of multiple organ failure following septic shock. *Am J Surg* 171: 221, 1996.
141. Manikis P, Jankowski S, Zhang H, et al: Correlation of serial blood lactate levels to organ failure and mortality after trauma. *Am J Emerg Med* 13: 619, 1995.
142. Nguyen HB, Rivers EP, Knoblich BP, et al: Early lactate clearance is associated with improved outcome in severe sepsis and septic shock. *Crit Care Med* 32: 1637, 2004.
143. Arnold RC, Shapiro NI, Jones AE, et al: Multi-center study of early lactate clearance as a determinant of survival in patients with presumed sepsis. *Shock* 32: 35, 2009.
144. Lima A, Bakker J: Noninvasive monitoring of peripheral perfusion. *Intensive Care Med* 31: 1316, 2005.
145. Cohn SM, Crookes BA, Proctor KG: Near-infrared spectroscopy in resuscitation. *J Trauma* 54: S199, 2003.
146. Mulier KE, Skarda DE, Taylor JH, et al: Near-infrared spectroscopy in patients with severe sepsis: correlation with invasive hemodynamic measurements. *Surg Infect* 9: 515, 2008.
147. Mesquida J, Masip J, Gili G, et al: Thenar oxygen saturation measured by near infrared spectroscopy as a noninvasive predictor of low central venous oxygen saturation in septic patients. *Intensive Care Med* 35: 1106, 2009.
148. Rhee P, Langdale L, Mock C, Gentilello LM: Near-infrared spectroscopy: continuous measurement of cytochrome oxidation during hemorrhagic shock. *Crit Care Med* 25: 166, 1997.
149. Ikossi DG, Knudson MM, Morabito DJ, et al: Continuous muscle tissue oxygenation in critically injured patients: a prospective observational study. *J Trauma* 61: 780, 2006.
150. Cohn SM, Nathens AB, Moore FA, et al: Tissue oxygen saturation predicts the development of organ dysfunction during traumatic shock resuscitation. *J Trauma* 62: 44, 2007.
151. Gomez H, Torres A, Polanco P, et al: Use of non-invasive NIRS during a vascular occlusion test to assess dynamic tissue O<sub>2</sub> saturation response. *Intensive Care Med* 34: 1600, 2008.
152. Wassenaar EB, Van den Brand JG: Reliability of near-infrared spectroscopy in people with dark skin pigmentation. *J Clin Monit Comput* 19: 195, 2005.
153. Gunn SR, Fink MP, Wallace B: Equipment review: the success of early goal-directed therapy for septic shock prompts evaluation of current approaches for monitoring the adequacy of resuscitation. *Crit Care* 9: 349, 2005.