

78 Upper Gastrointestinal Bleeding



REFERENCES

1. Van Leeradam ME, Vreeburg EM, Rauws AJ, et al: Acute upper GI bleeding: did anything change? *Am J Gastroenterol* 98: 1494, 2003.
2. Rockall TA, Logan RF, Devlin HB, et al: Incidence of and mortality from acute upper gastrointestinal hemorrhage in the United Kingdom: Steering Committee and members of the National Audit of Acute Upper Gastrointestinal Haemorrhage. *BMJ* 311: 222, 1995.
3. Wilcox CM, Alexander LN, Cotsonis G: A prospective characterization of upper gastrointestinal hemorrhage presenting with hematochezia. *Am J Gastroenterol* 92: 231, 1997.
4. Witting MD, Lagder L, Heins AE, et al: ED predictors of upper gastrointestinal track bleeding in patients without hematemesis. *Am J Emerg Med* 24: 280, 2006.
5. Barkun A, Bardou M, Marshall JK: Consensus recommendations for managing patients with nonvariceal upper gastrointestinal bleeding. *Ann Intern Med* 139: 843, 2003.
6. Gallagher EJ: Nasogastric tubes: hard to swallow. *Ann Emerg Med* 44: 138, 2004.
7. Witting MD, Lagder L, Heins AE, et al: Usefulness and validity of diagnostic nasogastric aspiration in patients without hematemesis. *Ann Emerg Med* 43: 525, 2004.
8. Ferguson JW, Tripathi D, Hayes PC: Review article: the management of acute variceal bleeding. *Aliment Pharmacol Ther* 18: 253, 2003.
9. Leontiadis GI, Sharma VK, Howden CW: Systematic review and meta-analysis of proton pump inhibitor therapy in peptic ulcer bleeding. *BMJ* 330: 568, 2005.
10. Palmer K: Acute upper gastrointestinal hemorrhage. *Br Med Bull* 83: 307, 2007.
11. Bardou M, Toubouti Y, Benharberou-Brun D, et al: Meta-analysis: proton-pump inhibition in high-risk patients with acute peptic ulcer bleeding. *Aliment Pharmacol Ther* 21: 677, 2005.
12. Julapalli VR, Graham DY: Appropriate use of intravenous proton pump inhibitors in the management of bleeding peptic ulcer. *Dig Dis Sci* 50: 1185, 2005.
13. Santander C, Gravalos RG, Gomez-Cedenilla A, et al: Antimicrobial therapy for *Helicobacter pylori* infection versus long-term maintenance antisecretion treatment in the prevention of recurrent hemorrhage from peptic ulcer: prospective nonrandomized trial on 125 patients. *Am J Gastroenterol* 91: 1549, 1996.
14. Frossard JL, Spahr L, Queneau PE, et al: Erythromycin intravenous bolus infusion in acute upper gastrointestinal bleeding: a randomized controlled double-blind trial. *Gastroenterology* 123: 17, 2002.
15. Corley DA, Stefan AM, Wolf M, et al: Early indicators of prognosis in upper gastrointestinal hemorrhage. *Am J Gastroenterol* 93: 336, 1998.
16. Stanley AJ, Ashley D, Dalton HR, et al: Outpatient management of patients with low-risk upper gastrointestinal haemorrhage: multicentre validation and prospective evaluation. *Lancet* 373: 42, 2009.
17. Romagnuolo J, Barkun AN, Enns R, et al: Simple clinical predictors may obviate urgent endoscopy in selected patients with nonvariceal upper gastrointestinal tract bleeding. *Arch Intern Med* 167: 265, 2007.