

146 Septic Shock



REFERENCES

- Angus DC, Linde-Zwirble WT, Lidicker J, et al: Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated cost of care. *Crit Care Med* 29: 1303, 2001.
- Martin GS, Mannino DM, Eaton S, et al: The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med* 348: 1546, 2003.
- Wang HE, Shapiro NI, Angus DC, et al: National estimates of severe sepsis in United States emergency departments. *Crit Care Med* 35: 1928, 2007.
- Watson RS, Carcillo JA, Linde-Zwirble WT, et al: The epidemiology of severe sepsis in children in the United States. *Am J Respir Crit Care Med* 167: 695, 2003.
- Bone RC, Balk RA, Cerra FB, et al: Definitions for organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. *Chest* 101: 1644, 1992.
- Levy MM, Fink MP, Marshall JC, et al: 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med* 31: 1250, 2003.
- Graciano AL, Balko JA, Rahn DS, et al: The Pediatric Multiple Organ Dysfunction Score (P-MODS): development and validation of an objective scale to measure the severity of multiple organ dysfunction in critically ill children. *Crit Care Med* 33: 1484, 2005.
- Goldstein B, Giroir B, Randolph A: International pediatric sepsis consensus conference: definitions for sepsis and organ dysfunction in pediatrics. *Pediatr Crit Care Med* 6: 2, 2005.
- Rangel-Frausto NS, Pittet D, Costigan M, et al: The natural history of the systemic inflammatory response syndrome (SIRS): a prospective study. *JAMA* 273: 117, 1995.
- Horn DL, Morrison DC, Opal SM, et al: What are the microbial components implicated in the pathogenesis of sepsis? Report on a symposium. *Clin Infect Dis* 31: 851, 2000.
- Hotchkiss RS, Karl IE: The pathophysiology and treatment of sepsis. *N Engl J Med* 348: 138, 2003.
- Remick DG: Pathophysiology of sepsis. *Am J Pathol* 170: 1435, 2007.
- Nguyen HB, Rivers EP, Knoblich BP, et al: Early lactate clearance is associated with improved outcome in severe sepsis and septic shock. *Crit Care Med* 32: 1637, 2004.
- Mikkelsen ME, Miltiades AN, Gaieski DF, et al: Serum lactate is associated with mortality in severe sepsis independent of organ failure and shock. *Crit Care Med* 37: 1670, 2009.
- Povoa P: C-reactive protein: a valuable marker of sepsis. *Intensive Care Med* 28: 235, 2002.
- Peres Bota D, Melot C, Lopes Ferreira F, et al: Infection Probability Score (IPS): a method to help assess the probability of infection in critically ill patients. *Crit Care Med* 31: 2579, 2003.
- Reinhart K, Karzai W, Meisner M: Procalcitonin as a marker of the systemic inflammatory response to infection. *Intensive Care Med* 26: 1193, 2000.
- Tugrul S, Esen F, Celebi S, et al: Reliability of procalcitonin as a severity marker in critically ill patients with inflammatory response. *Anesth Intensive Care* 30: 747, 2002.
- Guvan H, Altintop L, Baydin A, et al: Diagnostic value of procalcitonin levels as an early indicator of sepsis. *Am J Emerg Med* 20: 202, 2002.
- Ventetuolo CE, Levy MM: Biomarkers: diagnosis and risk assessment in sepsis. *Clin Chest Med* 29: 591, 2008.
- Sexton PM, Christopoulos G, Christopoulos A, et al: Procalcitonin has bioactivity at calcitonin receptor family complexes: potential mediator implications in sepsis. *Crit Care Med* 36: 1637, 2008.
- Uzzan B, Cohen R, Nicolas P, et al: Procalcitonin as a diagnostic test for sepsis in critically ill adults and after surgery or trauma: a systematic review and meta-analysis. *Crit Care Med* 34: 1996, 2006.
- Becker KL, Snider R, Nysten ES: Procalcitonin assay in systemic inflammation, infection, and sepsis: clinical utility and limitations. *Crit Care Med* 36: 941, 2008.
- Steinbach G, Rau B, Debarb AL, et al: Multicenter evaluation of a new immunoassay for procalcitonin measurement on the Kryptor system. *Clin Chem Lab Med* 42: 440, 2004.
- deWolf HK, Gunnewiek JK, Berk Y, et al: Comparison of a new procalcitonin assay from Roche with the established method on the Brahms Kryptor. *Clin Chem* 55: 1043, 2009.
- Russell JA: Management of sepsis. *N Engl J Med* 355: 1699, 2006.
- Rivers E, Nguyen B, Havstad S, et al: Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 345: 1368, 2001.
- Dellinger RP, Carlet JM, Masur H, et al: Surviving Sepsis Campaign guidelines for management of severe sepsis and septic shock. *Crit Care Med* 32: 858, 2004.
- Dellinger RP, Levy MM, Carlet JM, et al: Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock. *Crit Care Med* 36: 296, 2008.
- Brierley J, Carcillo JA, Choong K, et al: Clinical practice parameters for hemodynamic support of pediatric and neonatal septic shock: 2007 update from the American College of Critical Care Medicine. *Crit Care Med* 37: 666, 2009.
- Brower RG, Matthay MA, Morris A, et al: Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. The Acute Respiratory Distress Syndrome Network. *N Engl J Med* 342: 1301, 2000.
- Russell JA, Walley KR, Singer J, et al: Vasopressin versus norepinephrine infusion in patients with septic shock. *N Engl J Med* 358: 877, 2008.
- Carcillo JA, Fields AI: Clinical practice parameters for hemodynamic support of pediatric and neonatal patients in septic shock. *Crit Care Med* 30: 1365, 2002.
- Ceneviva G, Paschall JA, Maffei F, et al: Hemodynamic support in fluid-refractory pediatric septic shock. *Pediatrics* 102: e19, 1998.
- Hebert PC, Wells G, Blajchman MA, et al: A multicenter, randomized, controlled clinical trial of transfusion requirements in critical care. Transfusion Requirements in Critical Care Investigators, Canadian Critical Care Trials Group. *N Engl J Med* 340: 409, 1999.
- Sprung CL, Caralis PV, Marcial EH, et al: The effects of high-dose corticosteroids in patients with septic shock. A prospective, controlled study. *N Engl J Med* 311: 1137, 1984.
- Bone RC, Fisher CJ Jr, Clemmer TP, et al: A controlled clinical trial of high-dose methylprednisolone in the treatment of severe sepsis and septic shock. *N Engl J Med* 317: 653, 1987.
- Effect of high-dose glucocorticoid therapy on mortality in patients with clinical signs of systemic sepsis. The Veterans Administration Systemic Sepsis Cooperative Study Group. *N Engl J Med* 317: 659, 1987.
- Annane D, Sebille V, Charpentier C, et al: Effect of treatment with low doses of hydrocortisone and fludrocortisone on mortality in patients with septic shock. *JAMA* 288: 862, 2002.
- Sprung CL, Annane D, Keh D, et al: Hydrocortisone therapy for patients with septic shock. *N Engl J Med* 358: 111, 2008.
- Murray H, Marik PE: Etomidate for endotracheal intubation in sepsis: acknowledging the good while accepting the bad. *Chest* 127: 707, 2005.
- Kumar A, Roberts D, Wood KE, et al: Duration of hypotension before initiation of effective antimicrobial therapy is the critical determinant of survival in human septic shock. *Crit Care Med* 34: 1589, 2006.
- Bernard GR, Vincent JL, Laterre PF, et al: Efficacy and safety of recombinant human activated protein C for severe sepsis. *N Engl J Med* 344: 699, 2001.
- Vincent JL, Bernard GR, Beale R, et al: Drotrecogin alfa (activated) treatment in severe sepsis from the global open-label trial ENHANCE: further evidence for survival and safety and implications for early treatment. *Crit Care Med* 33: 2266, 2005.
- Abraham E, Laterre PF, Garg R, et al: Drotrecogin alfa (activated) for adults with severe sepsis and a low risk of death. *N Engl J Med* 353: 1332, 2005.
- Laterre PF, Abraham E, James JM, et al: ADDRESS (ADministration of DRotrecogin alfa [activated] in Early stage Severe Sepsis) long-term follow-up: one-year safety and efficacy evaluation. *Crit Care Med* 35: 1457, 2007.
- Nadel S, Goldstein B, Williams MD, et al: Drotrecogin alfa (activated) in children with severe sepsis: a multicentre phase III randomized controlled trial. *Lancet* 369: 836, 2007.
- Van Den Berghe G, Wouters P, Weekers F, et al: Intensive insulin therapy in the critically ill patients. *N Engl J Med* 345: 1359, 2001.
- Van den Berghe G, Wilmer A, Hermans G, et al: Intensive insulin therapy in the medical ICU. *N Engl J Med* 354: 449, 2006.
- Brunkhorst FM, Engel C, Bloos F, et al: Intensive insulin therapy and pentastarch resuscitation in severe sepsis. *N Engl J Med* 358: 125, 2008.
- Preiser JC, Devos P, Ruiz-Santana S, et al: A prospective randomised multi-centre controlled trial on tight glucose control by intensive insulin therapy in adult intensive care units: the GluControl study. *Intensive Care Med* 35(10): 1738, 2009.
- Finfer S, Chittock DR, Su SY, et al: Intensive versus conventional glucose control in critically ill patients. *N Engl J Med* 360: 1283, 2009.
- Werdan K, Pilz G, Bujdosó O, et al: Score-based immunoglobulin G therapy of patients with sepsis: the SBITS study. *Crit Care Med* 35: 2693, 2007.
- Pildal J, Gotzsche PC: Polyclonal immunoglobulin for treatment of bacterial sepsis: a systematic review. *Clin Infect Dis* 39: 38, 2004.
- Norrby-Teglund A, Haque KN, Hammarstrom L: Intravenous polyclonal IgM-enriched immunoglobulin therapy in sepsis: a review of clinical efficacy in relation to microbiological aetiology and severity of sepsis. *J Intern Med* 260: 509, 2006.
- Laupland KB, Kirkpatrick AW, Delaney A: Polyclonal intravenous immunoglobulin for the treatment of severe sepsis and septic shock in critically ill adults: a systematic review and meta-analysis. *Crit Care Med* 35: 2686, 2007.
- El-Nawawy A, El-Kinany H, Hamdy El-Sayed M, et al: Intravenous polyclonal immunoglobulin administration to sepsis syndrome patients: a prospective study in a pediatric intensive care unit. *J Trop Pediatr* 51: 271, 2005.
- Weinstein MP: Blood culture contamination: Persisting problems and partial progress. *J Clin Micro* 41: 2275, 2003.
- Reimer LG, Wilson KI, Weinstein MP: Update on detection of bacteremia and fungemia. *Clin Micro Rev* 10: 444, 1997.
- Weinstein M P, Towns ML, Quarley SM, et al: The clinical significance of positive blood cultures in the 1990s: a prospective comprehensive evaluation of the microbiology, epidemiology, and outcome of bacteremia and fungemia in adults. *Clin Infect Dis* 24: 584, 1997.
- Laupland KB, Church DL, Gregson DB: Blood cultures in ambulatory outpatients. *BMC Infect Dis* 5: 35, 2005.
- Stalnikowicz R, Block C: The yield of blood cultures in a department of emergency medicine. *Euro J Emerg Med* 8: 93, 2001.
- Kornberg AE, Jain N, Dannenhoffer R: Evaluation of false positive blood cultures: guidelines for early detection of contaminated cultures in febrile children. *Pediatr Emerg Care* 10: 20, 1994.
- McGowan KL, Foster JA, Coffin SE: Outpatient pediatric blood cultures: Time to positivity. *Pediatrics* 106: 251, 2000.
- Beckman SE, Diekema DJ, Doern GV: Determining the clinical significance of coagulase-negative staphylococci isolated from blood cultures. *Infect Control Hosp Epidemiol* 26: 559, 2005.
- Bates DW, Lee TH: Rapid classification of positive blood cultures. *JAMA* 267: 1962, 1992.

USEFUL WEB RESOURCES

Surviving Sepsis Campaign: International Guidelines for Management of Severe Sepsis and Septic Shock: 2008—http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2249616/pdf/134_2007_Article_934.pdf