

## 286 Anorexia Nervosa and Bulimia Nervosa

### REFERENCES

1. Hudson JI, Hiripi E, Pope HG Jr, Kessler RC: The prevalence and correlates of eating disorders in the national comorbidity survey replication. *Biol Psychiatry* 61: 348, 2007.
2. Mitchell JE, Crosby RD, Wonderlich SA, et al: Latent profile analysis of a cohort of patients with eating disorders not otherwise specified. *Int J Eat Disord* 40: S95, 2007.
3. Eaton EK, Kann L, Kinchen S, et al: Youth risk behavior surveillance—United States, 2005. *MMWR Surveill Summ* 55: 1, 2006.
4. Bulik CM, Tozzi F: Contemporary thinking about the role of genes and environment in eating disorders. *Epidemiol Psychiatr Soc* 13: 91, 2004.
5. McKnight Investigators: Risk factors for the onset of eating disorders in adolescent girls: results of the McKnight Longitudinal Risk Factor Study. *Am J Psychiatry* 160: 248, 2003.
6. Nattiv A, Agostini R, Drinkwater B, Yeager KK: The female athlete triad: the interrelatedness of disordered eating, amenorrhea, and osteoporosis. *Clin Sportsmed* 13: 405, 1994.
7. Herzog DB, Nussbaum KM, Marmor AK: Comorbidity and outcome in eating disorders. *Psych Clin North Am* 19: 843, 1996.
8. Pompili M, Girardi P, Tatarelli G, et al: Suicide and attempted suicide in eating disorders, obesity and weight-image concern. *Eat Behav* 7: 384, 2006.
9. Franko DL, Keel P: Suicidality in eating disorders: occurrence, correlates, and clinical implications. *Clin Psychol Rev* 26: 769, 2006.
10. Sullivan PF: Mortality in anorexia nervosa. *Am J Psychiatry* 152: 1073, 1995.
11. Patton GC: Mortality in eating disorders. *Psychol Med* 18: 947, 1998.
12. Lucas AR, Melton LJ, Crowson CS, O'Fallon WM: Long-term fracture risk among women with anorexia nervosa: a population-based cohort study. *Mayo Clinic Proc* 74: 972, 1999.
13. Brooks MJ, Melnik G: The refeeding syndrome: an approach to understanding its complications and preventing its occurrence. *Pharmacotherapy* 15: 713, 1995.
14. Fluoxetine in the treatment of bulimia nervosa. a multicenter, placebo-controlled, double-blind trial. Fluoxetine Bulimia Nervosa Collaborative Study Group. *Arch Gen Psychiatry* 49: 139, 1992.
15. Kaye WH, Nagata T, Weltzin TE, et al: Double-blind placebo-controlled administration of fluoxetine in restricting and restricting-purging-type anorexia nervosa. *Biol Psychiatry* 49: 644, 2001.
16. Steinhausen HC: The outcome of anorexia nervosa in the 20th century. *Am J Psychiatry* 159: 1284, 2002.
17. Keel PK, Mitchell JE, Miller KB, et al: Long-term outcome of bulimia nervosa. *Arch Gen Psychiatry* 56: 63, 1999.
18. Eckert ED, Halmi KA, Marchi P, et al: Ten-year follow-up of anorexia nervosa: clinical course and outcome. *Psychol Med* 25: 143, 1995.
19. Micali N, Simonoff E, Trasare J: Risk of major adverse perinatal outcomes in women with eating disorders. *Br J Psychiatry* 190: 255, 2007.

### ■ USEFUL WEB RESOURCES

National Eating Disorders Association—<http://www.nationaleatingdisorders.org>

National Association of Anorexia Nervosa and Associated Disorders—<http://www.anad.org>

The Academy for Eating Disorders. Includes free access to the International Journal of Eating Disorders—<http://www.aedweb.org>