

## REFERENCES

- McCaig LF, Ly N: National hospital ambulatory medical care survey: 2000 emergency department summary. *Adv Data* 326: 1, 2002. [PMID: 12661587]
- Nawar EW, Niska RW, Xu J: National Hospital Ambulatory Medical Care Survey: 2005 emergency department summary. *Adv Data* 386: 1, 2007. [PMID: 17703794]
- Christaki E, Opal SM: Is the mortality rate for septic shock really decreasing? *Curr Opin Crit Care* 14: 580, 2008. [PMID: 18787453]
- Khalid L, Dhakam SH: A review of cardiogenic shock in acute myocardial infarction. *Curr Cardiol Rev* 4: 34, 2008. [PMID: 19924275]
- Kumar A, Parillo JE: *Shock: Classification, Pathophysiology, and Approach to Management in Critical Care Medicine: Principles of Diagnosis and Management in the Adult*, 3rd ed. Philadelphia, PA: Mosby Inc.; 2008:379.
- Holt BD, Walsh RA: Normal physiology of the cardiovascular system, in Fuster V, Walsh RA, Harrington RA: *Hurst's the Heart*, 13th ed. New York: McGraw Hill; 2011.
- Shapiro NI, Howell MD, Talmor D, et al: Serum lactate as a predictor of mortality in emergency department patients with infection. *Ann Emerg Med* 45: 524, 2005. [PMID: 15855951]
- Bonanno FG: Hemorrhagic shock: the "physiology approach." *J Emerg Trauma Shock* 5: 285, 2012. [PMID: 23248495]
- Permpikul C, Sringam P, Tongyoo S: Therapeutic goal achievements during severe sepsis and septic shock resuscitation and their association with patients' outcomes. *J Med Assoc Thai* 97 (Suppl 3):S176, 2014.
- Arntfield RT, Millington SJ: Point of care cardiac ultrasound applications in the emergency department and intensive care unit: a review. *Curr Cardiol Rev* 8: 98, 2012. [PMID: 22894759]
- Atkinson PRT, McAuley DJ, Kendall RJ, et al: Abdominal and Cardiac Evaluation with Sonography in Shock (ACES): an approach by emergency physicians for the use of ultrasound in patients with undifferentiated hypotension. *Emerg Med J* 26: 87, 2009. [PMID: 19164614]
- Seif D, Perera P, Mailhot T, et al: Bedside ultrasound in resuscitation and the rapid ultrasound in shock protocol. *Crit Care Res Pract* 2012: 503254, 2012. [PMID: 23133747]
- Birkhahn RH, Gaeta TJ, Terry D, et al: Shock index in diagnosing early acute hypovolemia. *Am J Emerg Med* 23: 323, 2005. [PMID: 15915406]
- Karakitsos D, Labropoulos N, DeGroot E, et al: Real-time ultrasound-guided catheterisation of the internal jugular vein: a prospective comparison with the landmark technique in critical care patients. *Crit Care* 10: R162, 2006. [PMID: 17112371]
- Institute of Medicine: IOM report: the future of emergency care in the United States health system. *Acad Emerg Med* 13: 1081, 2006. [PMID: 17015421]
- Rivers E, Nguyen B, Havstad S, et al: Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 345: 1368, 2001. [PMID: 11794169]
- Dellinger RP, Levy MM, Carlet JM, et al: Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2008. *Crit Care Med* 36: 296, 2008. [PMID: 18158437]
- Sebat F, Musthafa AA, Johnson D, et al: Effect of a rapid response system for patients in shock on time to treatment and mortality during 5 years. *Crit Care Med* 35: 2568, 2007. [PMID: 17901831]
- ARISE Investigators, ANZICS Clinical Trials Group: Goal-directed resuscitation for patients with early septic shock. *N Engl J Med* 371: 1496, 2014. [PMID: 25272316]
- The ProCESS Investigators: A randomized trial of protocol-based care for early septic shock. *N Engl J Med* 370: 1683, 2014. [PMID: 24635773]
- Balas MC, Vasilevskis EE, Burke WJ, et al: Critical care nurses' role in implementing the "ABCDE bundle" into practice. *Crit Care Nurse* 32: 35, 2012. [PMID: 22467611]
- Foral JM, Voillet F, Pulina D, et al: Ventilator-associated pneumonia and ICU mortality in severe ARDS patients ventilated according to a lung-protective strategy. *Crit Care* 16: R65, 2012. [PMID: 22524447]
- Monnet X, Rienzo M, Osman D, et al: Passive leg raising predicts fluid responsiveness in critically ill patients. *Crit Care Med* 34: 1402, 2006. [PMID: 16540963]
- Anname D, Siami S, Jaber S, et al: Effects of fluid resuscitation with colloids vs crystalloids on mortality in critically ill patients presenting with hypovolemic shock: the CRISTAL randomized trial. *JAMA* 310: 1809, 2013. [PMID: 24108515]
- Dellinger RP, Levy MM, Rhodes A, et al: Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock, 2012. *Intensive Care Med* 39: 165, 2013. [PMID: 23361625]
- Yunos NM, Bellomo R, Hegarty C, Story D, Ho L, Bailey M: Association between a chloride-liberal vs chloride-restrictive intravenous fluid administration strategy and kidney injury in critically ill adults. *JAMA* 308: 1566, 2012. [PMID: 23073953]
- Shaw AD, Bagshaw SM, Goldstein SL, et al: Major complications, mortality, and resource utilization after open abdominal surgery: 0.9% saline compared to Plasma-Lyte. *Ann Surg* 255: 821, 2012. [PMID: 22470070]
- Asfar P, Meziani F, Hamel JF, et al: High versus low blood-pressure target in patients with septic shock. *N Engl J Med* 370: 1583, 2014. [PMID: 24635770]
- Russell JA: Is there a good MAP for septic shock? *N Engl J Med* 370: 1649, 2014. [PMID: 24635771]
- Cuschieri J, Rivers EP, Donnino MW, et al: Central venous-arterial carbon dioxide difference as an indicator of cardiac index. *Intensive Care Med* 31: 818, 2005. [PMID: 15803301]
- Weil MH, Nakagawa Y, Tang W, et al: Sublingual capnometry: a new noninvasive measurement for diagnosis and quantitation of severity of circulatory shock. *Crit Care Med* 27: 1225, 1999. [PMID: 10446813]
- Marik PE: Regional carbon dioxide monitoring to assess the adequacy of tissue perfusion. *Curr Opin Crit Care* 11: 245, 2005. [PMID: 15928474]
- Skarda DE, Mulier KE, Myers DE, et al: Dynamic near-infrared spectroscopy measurements in patients with severe sepsis. *Shock* 27: 348, 2007. [PMID: 17414414]
- Verdant C, De Backer D: How monitoring of the microcirculation may help us at the bedside. *Curr Opin Crit Care* 11: 240, 2005. [PMID: 15928473]
- Pinsky MR, Payen D: Functional hemodynamic monitoring. *Crit Care* 9: 566, 2005. [PMID: 16356240]
- Singer AJ, Thode HC Jr, Viccellio P, Pines JM: The association between length of emergency department boarding and mortality. *Acad Emerg Med* 18: 1324, 2011. [PMID: 22168198]
- Jones SE, Stiell IG, Nesbitt LP, et al: Nontraumatic out-of-hospital hypotension predicts in-hospital mortality. *Ann Emerg Med* 43: 106, 2004. [PMID: 14707949]
- Vincent JL, Gerlach H: Fluid resuscitation in severe sepsis and septic shock: an evidence-based review. *Crit Care Med* 32: S451, 2004. [PMID: 15542955]
- Caironi P, Tognoni G, Masson S, et al: Albumin replacement in patients with severe sepsis or septic shock. *N Engl J Med* 370: 1412, 2014. [PMID: 24635772]
- Finfer S, Bellomo R, Boyce N, et al: A comparison of albumin and saline for fluid resuscitation in the intensive care unit. *N Engl J Med* 350: 2247, 2004. [PMID: 15163774]
- American Thoracic Society: Evidence-based colloid use in the critically ill: American Thoracic Society Consensus Statement. *Am J Respir Crit Care Med* 170: 1247, 2004. [PMID: 15563641]
- Wiedermann CJ: Systematic review of randomized clinical trials on the use of hydroxyethyl starch for fluid management in sepsis. *BMC Emerg Med* 8: 1, 2008. [PMID: 18218122]
- Brunkhorst FM, Engel C, Bloos F, et al: Intensive insulin therapy and pentastarch resuscitation in severe sepsis. *N Engl J Med* 358: 125, 2008. [PMID: 18184958]
- Rivers EP, Kruse JA, Jacobsen G, et al: The influence of early hemodynamic optimization on biomarker patterns of severe sepsis and septic shock. *Crit Care Med* 35: 2016, 2007. [PMID: 17855815]
- Nguyen HB, Banta JE, Cho TW, et al: Mortality predictions using current physiologic scoring systems in patients meeting criteria for early goal-directed therapy and the severe sepsis resuscitation bundle. *Shock* 30: 23, 2008. [PMID: 18323748]

## USEFUL WEB RESOURCES

Critical Care Archives—<http://ccforum.com>  
 Critical Care Medicine Tutorials (SHOCK)—<http://ccmtutorials.com>, <http://ccmtutorials.com/cvs/index.htm>