

Hemodynamic Monitoring

H. Bryant Nguyen

David T. Huang

Michael R. Pinsky

REFERENCES

1. Vincent JL, Rhodes A, Perel A, et al: Clinical review: update on hemodynamic monitoring—a consensus of 16. *Crit Care* 15: 229, 2011. [PMID: 21884645]
2. Asfar P, Meziani F, Hamel JF, et al: High versus low blood-pressure target in patients with septic shock. *N Engl J Med* 370: 1583, 2014. [PMID: 24635770]
3. Bratton SL, Chestnut RM, Ghajar J, et al: Guidelines for the management of severe traumatic brain injury. I. Blood pressure and oxygenation. *J Neurotrauma* 24 (Suppl 1): S7, 2007. [PMID: NOT FOUND]
4. Bickell WH, Wall MJ, Jr, Pepe PE, et al: Immediate versus delayed fluid resuscitation for hypotensive patients with penetrating torso injuries. *N Engl J Med* 331: 1105, 1994. [PMID: 7935634]
5. Antonelli M, Levy M, Andrews PJ, et al: Hemodynamic monitoring in shock and implications for management. International Consensus Conference, Paris, France, 27–28 April 2006. *Intensive Care Med* 33: 575, 2007. [PMID: 17285286]
6. Pickering TG, Hall JE, Appel LJ, et al: Recommendations for blood pressure measurement in humans and experimental animals: part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Circulation* 111: 697, 2005. [PMID: 15699287]
7. Penaz J: Current photoelectric recording of blood flow through the finger. *Cesk Fysiol* 24: 349, 1975. [PMID: 1164745]
8. Lehman LW, Saeed M, Talmor D, Mark R, Malhotra A: Methods of blood pressure measurement in the ICU. *Crit Care Med* 41: 34, 2013. [PMID: 23269127]
9. Scheer B, Perel A, Pfeiffer UJ: Clinical review: complications and risk factors of peripheral arterial catheters used for haemodynamic monitoring in anaesthesia and intensive care medicine. *Crit Care* 6: 199, 2002. [PMID: 12133178]
10. Magder S: Central venous pressure: a useful but not so simple measurement. *Crit Care Med* 34: 2224, 2006. [PMID: 16763509]
11. Weil MH, Shubin H, Rosoff L: Fluid repletion in circulatory shock: central venous pressure and other practical guides. *JAMA* 192: 668, 1965. [PMID: 14280513]
12. Rivers E, Nguyen B, Havstad S, et al: Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 345: 1368, 2001. [PMID: 11794169]
13. Drazner MH, Rame JE, Stevenson LW, Dries DL: Prognostic importance of elevated jugular venous pressure and a third heart sound in patients with heart failure. *N Engl J Med* 345: 574, 2001. [PMID: 11529211]
14. Randazzo MR, Snoey ER, Levitt MA, Binder K: Accuracy of emergency physician assessment of left ventricular ejection fraction and central venous pressure using echocardiography. *Acad Emerg Med* 10: 973, 2003. [PMID: 12957982]
15. Lee BK, Lee HY, Jeung KW, Jung YH, Lee GS: Estimation of central venous pressure using inferior vena caval pressure from a femoral endovascular cooling catheter. *Am J Emerg Med* 31: 240, 2013. [PMID: 22980367]
16. Hadian M, Pinsky MR: Functional hemodynamic monitoring. *Curr Opin Crit Care* 13: 318, 2007. [PMID: 17468565]
17. Michard F, Boussat S, Chemla D, et al: Relation between respiratory changes in arterial pulse pressure and fluid responsiveness in septic patients with acute circulatory failure. *Am J Respir Crit Care Med* 162: 134, 2000. [PMID: 10903232]
18. Lopes MR, Oliveira MA, Pereira VO, Lemos IP, Auler JO, Jr, Michard F: Goal-directed fluid management based on pulse pressure variation monitoring during high-risk surgery: a pilot randomized controlled trial. *Crit Care* 11: R100, 2007. [PMID: 17822565]
19. Zhang Z, Lu B, Sheng X, Jin N: Accuracy of stroke volume variation in predicting fluid responsiveness: a systematic review and meta-analysis. *J Anesth* 25: 904, 2011. [PMID: 21892779]
20. Monnet X, Rienzo M, Osman D, et al: Passive leg raising predicts fluid responsiveness in the critically ill. *Crit Care Med* 34: 1402, 2006. [PMID: 16540963]
21. Summers RL, Shoemaker WC, Peacock WF, Ander DS, Coleman TG: Bench to bedside: electrophysiologic and clinical principles of noninvasive hemodynamic monitoring using impedance cardiography. *Acad Emerg Med* 10: 669, 2003. [PMID: 12782531]
22. Peacock WF, Summers RL, Vogel J, Emerman CE: Impact of impedance cardiography on diagnosis and therapy of emergent dyspnea: the ED-IMPACT trial. *Acad Emerg Med* 13: 365, 2006. [PMID: 16531605]
23. Kupersztych-Hagege E, Teboul JL, Artigas A, et al: Bioreactance is not reliable for estimating cardiac output and the effects of passive leg raising in critically ill patients. *Br J Anaesth* 111: 961, 2013. [PMID: 23985531]
24. Dark PM, Singer M: The validity of trans-esophageal Doppler ultrasonography as a measure of cardiac output in critically ill adults. *Intensive Care Med* 30: 2060, 2004. [PMID: 15368037]
25. Nguyen HB, Banta DP, Stewart G, et al: Cardiac index measurements by transcutaneous Doppler ultrasound and transthoracic echocardiography in adult and pediatric emergency patients. *J Clin Monit Comput* 24: 237, 2010. [PMID: 20563629]
26. Rajaram SS, Desai NK, Kalra A, et al: Pulmonary artery catheters for adult patients in intensive care. *Cochrane Database Syst Rev* 2: CD003408, 2013. [PMID: 23450539]
27. Reinhart K, Kuhn HJ, Hartog C, Bredle DL: Continuous central venous and pulmonary artery oxygen saturation monitoring in the critically ill. *Intensive Care Med* 30: 1572, 2004. [PMID: 15197435]
28. Dueck MH, Klimek M, Appenrodt S, Weigand C, Boerner U: Trends but not individual values of central venous oxygen saturation agree with mixed venous oxygen saturation during varying hemodynamic conditions. *Anesthesiology* 103: 249, 2005. [PMID: 16052106]
29. Trzeciak S, Dellinger RP, Abate NL, et al: Translating research to clinical practice: a 1-year experience with implementing early goal-directed therapy for septic shock in the emergency department. *Chest* 129: 225, 2006. [PMID: 16478835]
30. Shapiro NI, Howell MD, Talmor D, et al: Implementation and outcomes of the Multiple Urgent Sepsis Therapies (MUST) protocol. *Crit Care Med* 34: 1025, 2006. [PMID: 16484890]
31. Nguyen HB, Lynch EL, Mou JA, Lyon K, Wittlake WA, Corbett SW: The utility of a quality improvement bundle in bridging the gap between research and standard care in the management of severe sepsis and septic shock in the emergency department. *Acad Emerg Med* 14: 1079, 2007. [PMID: 17923718]
32. Jones AE, Focht A, Horton JM, Kline JA: Prospective external validation of the clinical effectiveness of an emergency department-based early goal-directed therapy protocol for severe sepsis and septic shock. *Chest* 132: 425, 2007. [PMID: 17573521]
33. A randomized trial of protocol-based care for early septic shock. *N Engl J Med* 370: 1683, 2014. [PMID: 24635773]
34. Kompanje EJ, Jansen TC, van der Hoven B, Bakker J: The first demonstration of lactic acid in human blood in shock by Johann Joseph Scherer (1814–1869) in January 1843. *Intensive Care Med* 33: 1967, 2007. [PMID: 17661014]
35. Fall PJ, Szerlip HM: Lactic acidosis: from sour milk to septic shock. *J Intensive Care Med* 20: 255, 2005. [PMID: 16145217]
36. Kruse JA, Zaidi SA, Carlson RW: Significance of blood lactate levels in critically ill patients with liver disease. *Am J Med* 83: 77, 1987. [PMID: 3605185]
37. Gallagher EJ, Rodriguez K, Touger M: Agreement between peripheral venous and arterial lactate levels. *Ann Emerg Med* 29: 479, 1997. [PMID: 9095008]
38. Bakker J, Gris P, Coffernils M, Kahn RJ, Vincent JL: Serial blood lactate levels can predict the development of multiple organ failure following septic shock. *Am J Surg* 171: 221, 1996. [PMID: 8619454]
39. Nguyen HB, Rivers EP, Knoblich BP, et al: Early lactate clearance is associated with improved outcome in severe sepsis and septic shock. *Crit Care Med* 32: 1637, 2004. [PMID: 15286537]
40. Jones AE, Shapiro NI, Trzeciak S, Arnold RC, Claremont HA, Kline JA: Lactate clearance vs central venous oxygen saturation as goals of early sepsis therapy: a randomized clinical trial. *JAMA* 303: 739, 2010. [PMID: 20179283]
41. Nguyen HB, Kuan WS, Batech M, et al: Outcome effectiveness of the severe sepsis resuscitation bundle with addition of lactate clearance as a bundle item: a multi-national evaluation. *Crit Care* 15: R229, 2011. [PMID: 21951322]
42. Mesquida J, Masip J, Gilí G, Artigas A, Baigorri F: Thunar oxygen saturation measured by near infrared spectroscopy as a noninvasive predictor of low central venous oxygen saturation in septic patients. *Intensive Care Med* 35: 1106, 2009. [PMID: 19183952]
43. Cohn SM, Nathens AB, Moore FA, et al: Tissue oxygen saturation predicts the development of organ dysfunction during traumatic shock resuscitation. *J Trauma* 62: 44, 2007. [PMID: 17215732]
44. Gomez H, Torres A, Polanco P, et al: Use of non-invasive NIRS during a vascular occlusion test to assess dynamic tissue O₂ saturation response. *Intensive Care Med* 34: 1600, 2008. [PMID: 18523754]
45. Guyette FX, Gomez H, Suffoletto B, et al: Prehospital dynamic tissue oxygen saturation response predicts in-hospital lifesaving interventions in trauma patients. *J Trauma Acute Care Surg* 72: 930, 2012. [PMID: 22491607]