

## REFERENCES

1. Mangili G, Aletti G, Frigerio L, et al: Palliative care for intestinal obstruction in recurrent ovarian cancer: a multivariate analysis. *Int J Gynecol Cancer* 15: 830, 2005. [PMID: 16174232]
2. Ripamonti C, Panzeri C, Groff L, et al: The role of somatostatin and octreotide in bowel obstruction: pre-clinical and clinical results. *Tumori* 87: 1, 2001. [PMID: 11669548]
3. Nelson R, Tsu B, Edwards S: Systematic review of prophylactic nasogastric decompression after abdominal operations. *Br J Surg* 92: 673, 2005. [PMID: 15912492]
4. Inman BA, Harel F, Tiguert R, et al: Routine nasogastric tubes are not required following cystectomy with urinary diversion: a comparative analysis of 430 patients. *J Urol* 170: 1888, 2003. [PMID: 14532800]
5. Witting MD, Magder L, Heins AE, et al: Emergency department predictors of upper gastrointestinal bleeding in patients presenting without hematemesis. *Am J Emerg Med* 24: 280, 2006. [PMID: 16635697]
6. Anderson RS, Witting MD, Pitera A, et al: Nasogastric aspiration in gastrointestinal hemorrhage. *Ann Emerg Med* 55: 364, 2010. [PMID: 20031264]
7. Witting MD, Magder L, Heins AE, et al: Usefulness and validity of diagnostic nasogastric aspiration in patients without hematemesis. *Ann Emerg Med* 43: 525, 2004. [PMID: 15039700]
8. Palamidessi N, Sinert R, Falzon L, et al: Nasogastric aspiration and lavage in emergency department patients with hematochezia or melena without hematemesis. *Acad Emerg Med* 17: 126, 2010. [PMID: 20370741]
9. Wilcox CM, Alexander LN, Cotsos G: A prospective characterization of upper gastrointestinal hemorrhage presenting with hematochezia. *Am J Gastroenterol* 92: 231, 1997. [PMID: 9040197]
10. Thomsen TW, Shaffer RW, Setnik GS: Videos in clinical medicine. Nasogastric intubation. *N Engl J Med* 354: e16, 2006. [PMID: 16641390]
11. Cullen L, Taylor D, Taylor S, Chu K: Nebulized lidocaine decreases the discomfort of nasogastric tube insertion: a randomized double-blind trial. *Ann Emerg Med* 44: 131, 2004. [PMID: 15278085]
12. Ozecelik DN, Daraca MA, Sivri B: Effectiveness of pre-emptive metoclopramide infusion in alleviating pain, discomfort, and nausea associated with nasogastric tube insertion: a randomized, double-blind, placebo-controlled trial. *Int J Clin Pract* 59: 1422, 2005. [PMID: 16351674]
13. McCullough S, Halton T, Mowbray D, Macfarland PI: Lingual sucrose reduces the pain response to nasogastric tube insertion: a randomized clinical trial. *Arch Dis Child Neonatal Ed* 93: F100, 2008. [PMID: 17634178]
14. Larkin GL, Claassen C: Trends in emergency department use of gastric lavage for poisoning events in the United States, 1993–2003. *Clin Toxicol* 45: 164, 2007. [PMID: 17364634]
15. Chinnock B, Afarian H, Minnigan H, et al: Physician clinical impression does not rule out spontaneous bacterial peritonitis in patients undergoing emergency department paracentesis. *Ann Emerg Med* 52: 268, 2008. [PMID: 18433932]
16. Wong CL, Holroyd-Leduc J, Thorpe KE, et al: Does this patient have bacterial peritonitis or portal hypertension? How do I perform a paracentesis and analyze the results? *JAMA* 299: 1166, 2008. [PMID: 18334692]
17. Mercaldi CJ, Lanes SF: Ultrasound guidance decreases complications and improves the cost of care among patients undergoing thoracentesis and paracentesis. *Chest* 143: 532, 2013. [PMID: 23381318]
18. Wollman B, D'Agostino HB: Percutaneous radiologic and endoscope gastrostomy a 3-year institutional analysis of procedure performance. *AJR Am J Roentgenol* 169: 1551, 1997. [PMID: 9393163]
19. Buscaglia JM: Common issues in PEG tubes—what every fellow should know. *Gastrointest Endosc* 64: 970, 2006. [PMID: 17140906]
20. Stumpf JL, Kurian RM, Vuong J, et al: Efficacy of a Creon delayed-release pancreatic enzyme protocol for clearing occluded feeding tubes. *Ann Pharmacother* 48: 483, 2014. [PMID: 24436458]
21. Korula J, Harma CA: Simple and inexpensive method of removal or replacement of gastrostomy tubes. *JAMA* 265: 1426, 1991. [PMID: 1999884]
22. Chait PG, Weinberg J, Connolly BL, et al: Retrograde percutaneous gastrostomy and gastrojejunostomy in 505 children. *Radiology* 201: 691, 1996. [PMID: 8939217]
23. Yaseen M, Steele MI, Grunow JE: Nonendoscopic removal of percutaneous endoscopic gastrostomy tubes: morbidity and mortality in children. *Gastrointest Endosc* 44: 235, 1996. [PMID: 8885339]
24. Erdil A, Genc H, Uygun A, et al: The buried bumper syndrome: the usefulness of retrieval PEG tubes in its management. *Turk J Gastroenterol* 19: 45, 2008. [PMID: 18386240]