

Approach to Nontraumatic Shock

Bret A. Nicks

John P. Gaillard

REFERENCES

- Holler JG, Bech CN, Henriksen DP, Mikkelsen S, Pedersen C, Lassen AT: Nontraumatic hypotension and shock in the emergency department and the prehospital setting, prevalence, etiology, and mortality: a systematic review. *PLoS One* 10: e0119331, 2015. [PMID: 25789927]
- Fleischmann C, Scherag A, Adhikari JKJ, et al: Assessment of global incidence and mortality of hospital-treated sepsis. Current estimates and limitations. *Am J Resp Crit Care Med* 193: 259, 2016. [PMID: 26414292]
- Jeger RV, Radovanovic D, Hunziker PR, et al: Ten-year trends in the incidence of cardiogenic shock. *Ann Intern Med* 149: 618, 2008. [PMID: 18981487]
- Thiele H, Zeymer U, Neumann FJ, et al: Intraaortic balloon support for myocardial infarction with cardiogenic shock. *N Engl J Med* 367: 1287, 2012. [PMID: 22920912]
- Guly HR, Bouamra O, Lecky FE: The incidence of neurogenic shock in patients with isolated spinal cord injury in the emergency department. *Resuscitation* 76:57, 2008. [PMID: 17688997]
- Russell JA, Rush B, Boyd J: Pathophysiology of septic shock. *Crit Care Clin* 34: 43, 2018. [PMID: 29149941]
- Biering-Sørensen F, Biering-Sørensen T, Liu N, et al: Alterations in cardiac autonomic control in spinal cord injury. *Auton Neurosci* 209: 4, 2018. [PMID: 28228335]
- Ashruf JF, Bruining HA, Ince C: New insights into the pathophysiology of cardiogenic shock: the role of the microcirculation. *Curr Opin Crit Care* 19: 381, 2013. [PMID: 23995131]
- Holler JG, Jensen HK, Henriksen DP, et al: Etiology of shock in the emergency department; a 12 year population based cohort study. *Shock* 51:60, 2019. [PMID: 27984523]
- Kheng CP, Rahman NH: The use of end-tidal carbon dioxide monitoring in patients with hypotension in the emergency department. *Int J Emerg Med* 5: 31, 2012. [PMID: 22828152]
- Holt BD, Walsh RA: Normal physiology of the cardiovascular system, in Fuster V, Walsh RA, Harrington RA (eds): *Hurst's the Heart*, 13th ed. New York, NY: McGraw Hill; 2011.
- Shapiro NI, Howell MD, Talmor D, et al: Serum lactate as a predictor of mortality in emergency department patients with infection. *Ann Emerg Med* 45: 524, 2005. [PMID: 15855951]
- Bonanno FG: Hemorrhagic shock: the "physiology approach." *J Emerg Trauma Shock* 5: 285, 2012. [PMID: 23248495]
- Oedorf K, Day D, Lior Y, et al: Serum lactate predicts adverse outcomes in emergency department patients with and without infection. *West J Emerg Med* 18: 258, 2017. [PMID: 28210362]
- Parenica J, Jarkovsky J, Malaska J, et al: Infectious complications and immune/inflammatory response in cardiogenic shock: a prospective observational study. *Shock* 47: 165, 2017. [PMID: 27749762]
- Singer M, Deutschman CS, Seymour CW, et al: The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 315: 801, 2016. [PMID: 26903338]
- Arntfield RT, Millington SJ: Point of care cardiac ultrasound applications in the emergency department and intensive care unit: a review. *Curr Cardiol Rev* 8: 98, 2012. [PMID: 22894759]
- Atkinson PRT, McAuley DJ, Kendall RJ, et al: Abdominal and Cardiac Evaluation with Sonography in Shock (ACES): an approach by emergency physicians for the use of ultrasound in patients with undifferentiated hypotension. *Emerg Med J* 26: 87, 2009. [PMID: 19164614]
- Seif D, Perera P, Mailhot T, et al: Bedside ultrasound in resuscitation and the rapid ultrasound in shock protocol. *Crit Care Res Pract* 2012: 503254, 2012. [PMID: 23133747]
- Mok KL: Make it SIMPLE: enhanced shock management by focused cardiac ultrasound. *J Intensive Care* 4: 51, 2016. [PMID: 27529027]
- Birkhahn RH, Gaeta TJ, Terry D, et al: Shock index in diagnosing early acute hypovolemia. *Am J Emerg Med* 23: 323, 2005. [PMID: 15915406]
- Karakitsos D, Labropoulos N, DeGroot E, et al: Real-time ultrasound-guided catheterization of the internal jugular vein: a prospective comparison with the landmark technique in critical care patients. *Crit Care* 10: R162, 2006. [PMID: 17112371]
- Institute of Medicine: IOM report: the future of emergency care in the United States health system. *Acad Emerg Med* 13: 1081, 2006. [PMID: 17015421]
- Rivers E, Nguyen B, Havstad S, et al: Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 345: 1368, 2001. [PMID: 11794169]
- Dellinger RP, Levy MM, Carlet JM, et al: Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2008. *Crit Care Med* 36: 296, 2008. [PMID: 18158437]
- Sebat F, Mustafa AA, Johnson D, et al: Effect of a rapid response system for patients in shock on time to treatment and mortality during 5 years. *Crit Care Med* 35: 2568, 2007. [PMID: 17901831]
- ARISE Investigators, ANZICS Clinical Trials Group: Goal-directed resuscitation for patients with early septic shock. *N Engl J Med* 371: 1496, 2014. [PMID: 25272316]
- The ProCESS Investigators: A randomized trial of protocol-based care for early septic shock. *N Engl J Med* 370: 1683, 2014. [PMID: 24635773]
- Balas MC, Vasilevskis EE, Burke WJ, et al: Critical care nurses' role in implementing the "ABCDE bundle" into practice. *Crit Care Nurse* 32: 35, 2012. [PMID: 22467611]
- Foral JM, Voillet F, Pulina D, et al: Ventilator-associated pneumonia and ICU mortality in severe ARDS patients ventilated according to a lung-protective strategy. *Crit Care* 16: R65, 2012. [PMID: 22524447]
- Monnet X, Rienzo M, Osman D, et al: Passive leg raising predicts fluid responsiveness in critically ill patients. *Crit Care Med* 34: 1402, 2006. [PMID: 16540963]
- Annane D, Siami S, Jaber S, et al: Effects of fluid resuscitation with colloids vs crystalloids on mortality in critically ill patients presenting with hypovolemic shock: the CRISTAL randomized trial. *JAMA* 310: 1809, 2013. [PMID: 24108515]
- Semler MW, Self WH, Wanderer JP, et al: Balanced crystalloids versus saline in critically ill adults. *N Engl J Med* 378: 829, 2018. [PMID: 29485925]
- Dellinger RP, Levy MM, Rhodes A, et al: Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock, 2012. *Intensive Care Med* 39: 165, 2013. [PMID: 23361625]
- Yunos NM, Bellomo R, Hegarty C, Story D, Ho L, Bailey M: Association between a chloride-liberal vs chloride-restrictive intravenous fluid administration strategy and kidney injury in critically ill adults. *JAMA* 308: 1566, 2012. [PMID: 23073953]
- Raghunathan K, Bonavia A, Nathanson BH, et al: Association between initial fluid choice and subsequent in-hospital mortality during the resuscitation of adults with septic shock. *Anesthesiology* 123: 1385, 2015. [PMID: 26414499]
- Asfar P, Meziani F, Hamel JF, et al: High versus low blood-pressure target in patients with septic shock. *N Engl J Med* 370: 1583, 2014. [PMID: 24635770]
- Russell JA: Is there a good MAP for septic shock? *N Engl J Med* 370: 1649, 2014. [PMID: 24635771]
- Cardenas-Garcia J, Schaub KF, Belchikov YG, et al: Safety of peripheral intravenous administration of vasoactive medication. *J Hosp Med* 10: 581, 2015. [PMID: 26014852]
- Loubani OM, Green RS: A systematic review of extravasation and local tissue injury from administration of vasopressors through peripheral intravenous catheters and central venous catheters. *J Crit Care* 30: 653.e9, 2015. [PMID: 25669592]
- Cuschieri J, Rivers EP, Donnino MW, et al: Central venous-arterial carbon dioxide difference as an indicator of cardiac index. *Intensive Care Med* 31: 818, 2005. [PMID: 15803301]
- Weil MH, Nakagawa Y, Tang W, et al: Sublingual capnometry: a new noninvasive measurement for diagnosis and quantitation of severity of circulatory shock. *Crit Care Med* 27: 1225, 1999. [PMID: 10446813]
- Marik PE: Regional carbon dioxide monitoring to assess the adequacy of tissue perfusion. *Curr Opin Crit Care* 11: 245, 2005. [PMID: 15928474]
- Skarda DE, Mulier KE, Myers DE, et al: Dynamic near-infrared spectroscopy measurements in patients with severe sepsis. *Shock* 27: 348, 2007. [PMID: 17414414]
- Verdant C, De Backer D: How monitoring of the microcirculation may help us at the bedside. *Curr Opin Crit Care* 11: 240, 2005. [PMID: 15928473]
- Pinsky MR, Payen D: Functional hemodynamic monitoring. *Crit Care* 9: 566, 2005. [PMID: 16356240]
- Singer AJ, Thode HC Jr, Viccellio P, Pines JM: The association between length of emergency department boarding and mortality. *Acad Emerg Med* 18: 1324, 2011. [PMID: 22168198]
- Jones SE, Stiell IG, Nesbitt LP, et al: Nontraumatic out-of-hospital hypotension predicts in-hospital mortality. *Ann Emerg Med* 43: 106, 2004. [PMID: 14707949]
- Vincent JL, Gerlach H: Fluid resuscitation in severe sepsis and septic shock: an evidence-based review. *Crit Care Med* 32: S451, 2004. [PMID: 15542955]
- Caironi P, Tognoni G, Masson S, et al: Albumin replacement in patients with severe sepsis or septic shock. *N Engl J Med* 370: 1412, 2014. [PMID: 24635772]
- Finfer S, Bellomo R, Boyce N, et al: A comparison of albumin and saline for fluid resuscitation in the intensive care unit. *N Engl J Med* 350: 2247, 2004. [PMID: 15163774]
- American Thoracic Society: Evidence-based colloid use in the critically ill: American Thoracic Society Consensus Statement. *Am J Respir Crit Care Med* 170: 1247, 2004. [PMID: 15563641]
- Wiedermann CJ: Systematic review of randomized clinical trials on the use of hydroxyethyl starch for fluid management in sepsis. *BMC Emerg Med* 8: 1, 2008. [PMID: 18218122]
- Brunkhorst FM, Engel C, Bloos F, et al: Intensive insulin therapy and pentastarch resuscitation in severe sepsis. *N Engl J Med* 358: 125, 2008. [PMID: 18184958]
- Self WH, Semler MW, Wanderer JP, et al: Balanced crystalloids versus saline in noncritically ill adults. *N Engl J Med* 378: 819, 2018. [PMID: 29485926]
- Semler MW, Self WH, Wanderer JP, et al: Balanced crystalloids versus saline in critically ill adults. *N Engl J Med* 378: 829, 2018. [PMID: 29485925]
- Rochwerg B, Millen T, Austin P, et al: Fluids in Sepsis and Septic Shock (FISSH): protocol for a pilot randomised controlled trial. *BMJ Open* 7: e017602, 2017. [PMID: 28729329]
- Serpa Neto A, Martin Loeches I, Klanderman RB, et al: Balanced versus isotonic saline resuscitation—a systematic review and meta-analysis of randomized controlled trials in operation rooms and intensive care units. *Ann Transl Med* 5: 323, 2017. [PMID: 28861420]
- Zayed YZM, Aburahma AMY, Barbarawi MO, et al: Balanced crystalloids versus isotonic saline in critically ill patients: systematic review and meta-analysis. *J Intensive Care* 6: 51, 2018. [PMID: 30140441]
- McIntyre L, Taljaard M, McArdle T, et al: FLUID trial: a protocol for a hospital-wide open-label cluster crossover pragmatic comparative effectiveness randomized pilot trial. *BMJ Open* 8: e022780, 2018. [PMID: 30139908]

61. Zampieri FG, Azevedo LCP, Corrêa TD, et al: Study protocol for the Balanced Solution versus Saline in Intensive Care Study (BaSICS): a factorial randomised trial. *Crit Care Resusc* 19: 175, 2017. [PMID: 28651514]
62. Friederich A, Martin N, Swanson MB, et al: Normal saline solution and lactated Ringer's solution have a similar effect on quality of recovery: a randomized controlled trial. *Ann Emerg Med* 2018, August 23. [Epub ahead of print] [PMID: 30146446]
63. Myburgh JA, Mythen MG: Resuscitation fluids. *N Engl J Med* 369: 1243, 2013. [PMID: 24066745]
64. Kimmoun A, Novy E, Auchet T, et al: Hemodynamic consequences of severe lactic acidosis in shock states: from bench to bedside. *Crit Care* 19: 175, 2015. [PMID: 25887061]
65. Kim HJ, Son YK, An WS: Effect of sodium bicarbonate administration on mortality in patients with lactic acidosis: a retrospective analysis. *PLoS One* 8: e65283, 2013. [PMID: 23755210]
66. Rivers EP, Kruse JA, Jacobsen G, et al: The influence of early hemodynamic optimization on biomarker patterns of severe sepsis and septic shock. *Crit Care Med* 35: 2016, 2007. [PMID: 17855815]
67. Nguyen HB, Banta JE, Cho TW, et al: Mortality predictions using current physiologic scoring systems in patients meeting criteria for early goal-directed therapy and the severe sepsis resuscitation bundle. *Shock* 30: 23, 2008. [PMID: 18323748]