

REFERENCES

1. Geeraedts LM, Kaasjager HA, van Vugt AB, Frolke JP: Exsanguination in trauma: a review of diagnostics and treatment options. *Injury* 40: 11, 2009. [PMID: 19135193]
2. <http://www.cdc.gov/hicpac/pdf/guidelines/bsi-guidelines-2011.pdf> (Centers for Disease Control and Prevention: Guidelines for Intravascular Catheter-Related Infections.) Accessed April 14, 2014.
3. Malach T, Jerassy Z, Rudensky B, et al: Prospective surveillance of phlebitis associated with peripheral intravenous catheters. *Am J Infect Control* 34: 308, 2006. [PMID: 16765211]
4. Zarate L, Mandelco B, Wishaw R, et al: Peripheral intravenous catheters started in pre-hospital and emergency department settings. *J Trauma Nurs* 15: 47, 2008. [PMID: 18690133]
5. Brannam L, Blaivas M, Lyon M, et al: Emergency nurses' utilization of ultrasound guidance for placement of peripheral intravenous lines in difficult-access patients. *Acad Emerg Med* 11: 1361, 2004. [PMID: 15576530]
6. Bauman M, Braude D, Crandall C: Ultrasound-guidance vs. standard technique in difficult vascular access patients by ED technicians. *Am J Emerg Med* 27: 135, 2009. [PMID: 19371518]
7. Werner SL, Jones RA, Emerman CL: Effect of hip abduction and external rotation on femoral vein exposure for possible cannulation. *J Emerg Med* 35: 73, 2008. [PMID: 17976787]
8. Stone MB, Price DD, Anderson BS: Investigation of the effect of reverse Trendelenburg on the cross-sectional area of the femoral vein. *J Emerg Med* 30: 211, 2006. [PMID: 16567260]
9. Shiloh AL, Savel RH, Paulin LM, et al: Ultrasound-guided catheterization of the radial artery: a systematic review and meta-analysis of randomized controlled trials. *Chest* 139: 524, 2011. [PMID: 20724734]