

20 Blood Gases

REFERENCES

1. Stein PD, Goldhaber SZ, Henry JW: Alveolar-arterial oxygen gradient in the assessment of acute pulmonary embolism. *Chest* 107: 139, 1995.
2. Bernard G, Artigas A, Brigham K, et al: The American-European Consensus Conference on ARDS: Definitions, mechanisms, relevant outcomes, and clinical trial coordination. *Am J Respir Crit Care Med* 149: 818, 1994.
3. Severinghaus JW, Honda Y: History of blood gas analysis. VII. Pulse oximetry. *J Clin Monit* 3: 135, 1987.
4. Barker SJ, Tremper KK: The effect of carbon monoxide inhalation on pulse oximetry and transcutaneous PO₂. *Anesthesiology* 66: 677, 1987.
5. Barker SJ, Tremper KK, Hyatt J: Effects of methemoglobinemia on pulse oximetry and mixed venous oximetry. *Anesthesiology* 70: 112, 1989.
6. Marik PE: Gastric intramucosal pH: a better predictor of multiorgan dysfunction syndrome and death than oxygen-derived variables in patients with sepsis. *Chest* 104: 225, 1993.
7. Doglio GR, Pusajo JF, Egurrola MA, et al: Gastric mucosal pH as a prognostic index of mortality in critically ill patients. *Crit Care Med* 19: 1037, 1991.
8. Marik PE: Sublingual capnography: a clinical validation study. *Chest* 120: 701, 1993.
9. Creteur J: Muscle StO₂ in critically ill patients. *Curr Opin Crit Care* 14: 361, 2008.
10. Wo CC, Shoemaker WC, Appel PL: Unreliability of blood pressure and heart rate to evaluate cardiac output in emergency resuscitation and critical illness. *Crit Care Med* 21: 218, 1993.
11. Rady MY, Rivers EP, Nowak RM: Resuscitation of the critically ill in the ED: responses of blood pressure, heart rate, shock index, central venous oxygen saturation, and lactate. *Am J Emerg Med* 14: 218, 1996.
12. Dries DJ, Waxman K: Adequate resuscitation of burn patients may not be measured by urine output and vital signs. *Crit Care Med* 19: 327, 1991.
13. Shapiro NI, Howell MD, Talmor D, et al: Serum lactate as a predictor of mortality in emergency department patients with infection. *Ann Emerg Med* 45: 524, 2005.
14. Nguyen HB, Rivers EP, Knoblich BP, et al: Early lactate clearance is associated with improved outcome in severe sepsis and septic shock. *Crit Care Med* 32: 1637, 2004.
15. Abramson D, Scalea TM, Hitchcock R, et al: Lactate clearance and survival following injury. *J Trauma* 35: 584, 1993.
16. Cohen RD, Woods HF: *Clinical and Biochemical Aspects of Lactic Acidosis*. Oxford, UK, Blackwell, 1976, pp. 40, 180.
17. Reinhart K, Rudolph T, Bredle DL: Comparison of central-venous to mixed-venous oxygen saturation during changes in oxygen supply/demand. *Chest* 95: 1216, 1989.
18. Scalea TM, Hartnett RW, Duncan AO, et al: Central venous oxygen saturation: a useful clinical tool in trauma patients. *J Trauma* 30: 1539, 1990.
19. Rivers E, Nguyen B, Havstad S, et al: Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 345: 1368, 2001.
20. Silvestri S, Ralls GA, Krauss B, et al: The effectiveness of out-of-hospital use of continuous end-tidal CO₂ monitoring on the rate of unrecognized misplaced intubation within a regional emergency medical services system. *Ann Emerg Med* 45: 497, 2005.
21. Falk J, Rackow E, Well M: End-tidal CO₂ concentration during cardiopulmonary resuscitation. *N Engl J Med* 318: 607, 1988.

■ USEFUL WEB RESOURCE

Altitude.org: A High Altitude Resource—<http://www.altitude.org>