

296 The Morbidly Obese Patient

REFERENCES

- National Center for Health Statistics, Centers for Disease Control and Prevention: Available at: <http://www.cdc.gov/nccdphp/dnpa/obesity/trend/index.htm>. Accessed 2008.
- Bertakis KD, Azari R: The influence of obesity, alcohol abuse, and smoking on utilization of health care services. *Fam Med* 38: 427, 2006.
- Hering E, Pritsker I, Gonchar L, Pillar G: Obesity in children is associated with increased health care use. *Clin Pediatr (Phila)* 48: 812, 2009.
- Platts-Mills TF, Burg MD, Snowden B: Obese patients with abdominal pain presenting to the emergency department do not require more time or resources for evaluation than nonobese patients. *Acad Emerg Med* 12: 778, 2005.
- Chen EH, Shofer FS, Hollander JE, et al: Emergency physicians do not use more resources to evaluate obese patients with acute abdominal pain. *Am J Emerg Med* 25: 925, 2007.
- Varon J, Marik P: Management of the obese critically ill patient. *Crit Care Clin* 17: 187, 2001.
- Joffe A, Wood K: Obesity in critical care. *Curr Opin Anesthesiol* 20: 113, 2007.
- Brown CV, Neville AL, Rhee P, et al: The impact of obesity on the outcomes of 1,153 critically injured blunt trauma patients. *J Trauma* 59: 1048, 2005.
- Kleiser C, Rosario AS, Mensink GBM, et al: Potential determinants of obesity among children and adolescents in Germany: results from cross-sectional KiGGS Study. *BMC Public Health* 9: 46, 2009.
- McGee DL: Body mass index and mortality: a meta-analysis based on person-level data from twenty-six observational studies. *Ann Epidemiol* 15: 87, 2005.
- Romero-Corral A, Montori VM, Somers VK, et al: Association of bodyweight with total mortality and with cardiovascular events in coronary artery disease: a systematic review of cohort studies. *Lancet* 368: 666, 2006.
- Rocha AT, de Vasconcellos AG, da Luz Neto ER, et al: Risk of venous thromboembolism and efficacy of thromboprophylaxis in hospitalized obese patients and in obese patients undergoing bariatric surgery. *Obes Surg* 16: 1645, 2006.
- Goldhaber S, Grodstein E, Stampfer M, et al: A prospective study of risk factors for pulmonary embolism in women. *JAMA* 277: 642, 1997.
- Taylor EN, Stampfer MJ, Curhan GC: Obesity, weight gain, and the risk of kidney stones. *JAMA* 293: 455, 2005.
- Calle EE, Rodriguez C, Walker-Thurmond K, Thun MJ: Overweight, obesity, and mortality from cancer in a prospectively studied cohort of U.S. adults. *N Engl J Med* 348: 1625, 2003.
- Petrella RJ, Lattanzio CN: Does counseling help patients get active? Systemic review of the literature. *Can Fam Physician* 48: 72, 2002.
- Alpert MA, Lambert CR, Panayiotou H, et al: Relation of duration of morbid obesity to left ventricular mass, systolic function, and diastolic filling, and effect of weight loss. *Am J Cardiol* 76: 1194, 1995.
- Lazarus R, Sparrow D, Weiss S: Effects of obesity and fat distribution on ventilatory function: the normative aging study. *Chest* 111: 891, 1997.
- Perilli V, Sollazzi L, Modesti C, et al: Comparison of positive end-expiratory pressure with reverse Trendelenburg position in morbidly obese patients undergoing bariatric surgery: effects on hemodynamics and pulmonary gas exchange. *Obes Surg* 13: 605, 2003.
- Goldhaber SZ, Tapson VF: A prospective registry of 5,451 patients with ultrasound confirmed deep vein thrombosis. *Am J Cardiol* 93: 259, 2004.
- Sharma OP, Owsanski MF, Joseph RJ, et al: Venous thromboembolism in trauma patients. *Ann Surg* 73: 1173, 2007.
- Metylwy M, Ong KJ, Ledger WL, Li TC: Does high body mass index increase the risk of miscarriage after spontaneous and assisted conception? A meta-analysis of the evidence. *Fertil Steril* 90: 714, 2008.
- Abenhaim HA, Kinch RA, Morin L, et al: Effect of prepregnancy body mass index category on obstetrical and neonatal outcomes. *Arch Gynecol Obstet* 275: 39, 2006.
- Muppala H, Rafi J, Arthur I: Morbidly obese woman unaware of pregnancy until full-term and complicated by intraamniotic sepsis with pseudomonas. *Infect Dis Obstet Gynecol* 51689, 2007.
- Xiang H, Smith GA, Wilkins JR III, et al: Obesity and risk of nonfatal unintentional injuries. *Am J Prev Med* 29: 41, 2005.
- Bostman OM: Body mass index of patients with elbow and ankle fractures requiring surgical treatment. *J Trauma* 37: 62, 1994.
- Bostman OM: Body weight related to loss of reduction of fractures of the distal tibia and ankle. *J Bone Joint Surg Br* 77: 101, 1995.
- Whitlock G, Norton R, Jackson R, et al: Is body mass index a risk factor for motor vehicle driver injury? A cohort study with prospective and retrospective outcomes. *Int J Epidemiol* 32: 147, 2003.
- Teran-Santos J, Jimenez-Gomez A, Cordero-Guevara J: The association between sleep apnea and the risk of traffic accidents. *N Engl J Med* 18: 340, 1999.
- Bochicchio GV, Joshi M, Bochicchio K, et al: Impact of obesity in the critically ill trauma patient: a prospective study. *J Am Coll Surg* 203: 533, 2006.
- Ryb GE, Dischinger PC: Injury severity and outcome of overweight and obese patients after vehicular trauma: a crash injury research and engineering network (CIREN) study. *J Trauma* 64: 406, 2008.
- Arbab S, Wahl WL, Hemmila MR, et al: The cushion effect. *J Trauma* 54: 1090, 2003.
- Brown CV, Neville AL, Salim A, et al: The impact of obesity on severely injured children and adolescents. *J Pediatr Surg* 41: 88, 2006.
- Zarzaa BL, Marshall SW: Motor vehicle crashes obesity and seat belt use: a deadly combination? *J Trauma* 64: 412, 2008.
- Luber SD, Fischer DR, Venkat A: Care of the bariatric surgery patient in the emergency department. *J Emerg Med* 34: 13, 2008.
- Edwards ED, Jacob BP, Gagner M, Pomp A: Presentation and management of common post-weight loss surgery problems in the emergency department. *Ann Emerg Med* 47: 160, 2006.
- Lemmens HJ, Brodsky JB: The dose of succinylcholine in morbid obesity. *Anesth Analg* 102: 438, 2006.
- Schwartz AE, Matteo RS, Ornstein E, et al: Pharmacokinetics and pharmacodynamics of vecuronium in the obese surgical patient. *Anesth Analg* 74: 515, 1992.
- Spruill WJ, Wade WE, Huckaby WG, Leslie RB: Achievement of anticoagulation by using a weight-based heparin dosing protocol for obese and nonobese patients. *Am J Health Syst Pharm* 59: 1788, 2002.
- Schwiesow SJ, Wessell AM, Steyer TE: Use of a modified dosing weight for heparin therapy in a morbidly obese patient. *Ann Pharmacol* 39: 753, 2005.
- Barras MA, Duffel SB, Atherton JJ, Green B: Individualized compared with conventional dosing of enoxaparin. *Clin Pharmacol Ther* 83: 882, 2007.
- Davidson BL, Buller HR, Decousus H, et al: Effect of obesity on outcomes after fondaparinux, enoxaparin, or heparin treatment for acute venous thromboembolism in the Matisse trials. *J Thromb Haemost* 5: 1191, 2007.
- Brodsky JB, Lemmens HJ: Regional anesthesia and obesity. *Obes Surg* 17: 1146, 2007.
- Casati A, Putzu M: Anesthesia in the obese patient: pharmacokinetic considerations. *J Clin Anesth* 17: 134, 2005.
- Venn RM, Hell J, Grounds RM: Respiratory effects of dexmedetomidine in the surgical patient requiring intensive care. *Crit Care* 4: 302, 2000.
- Bauer LA, Black DJ, Lill JS: Vancomycin dosing in morbidly obese patients. *Eur J Clin Pharmacol* 54: 621, 1998.
- Pai MP, Bearden DT: Antimicrobial dosing considerations in obese adult patients. *Pharmacotherapy* 27: 1081, 2007.
- Hollenstein UM, Brunner M, Schmid R, Muller M: Soft tissue concentrations of ciprofloxacin in obese and lean subjects following weight-adjusted dosing. *Int J Obes Relat Metab Disord* 25: 354, 2001.
- Berthoud MC, Peacock JE, Reilly CS: Effectiveness of preoxygenation in morbidly obese patients. *Br J Anaesth* 67: 464, 1991.
- Wieczorek PM, Schriker T, Vinet B, Backman SB: Airway topicalization in morbidly obese patients using atomized lidocaine: 2% compared with 4%. *Anaesthesia* 63: 316, 2007.
- Natalini G, Franceschetti MI, Pantelidi MT, et al: Comparison of the standard laryngeal mask airway and the ProSeal laryngeal mask airway in obese patients. *Br J Anaesth* 90: 323, 2003.
- Levitian RM, Ocroh EA, Stuart S, et al: Use of intubating laryngeal mask airway by medical and nonmedical personnel. *Am J Emerg Med* 18: 12, 2000.
- Fraley MA, Birchem JA, Senkottaiyan N, Alpert MA: Obesity and the electrocardiogram. *Obesity Reviews* 6: 275, 2005.
- Maxwell MH, Waks AU, Schrot PC, et al: Error in blood-pressure measurement due to incorrect cuff size in obese patients. *Lancet* 2: 33, 1982.
- Keyes LE, Frazee BW, Snoey ER, et al: Ultrasound-guided brachial and basilic vein cannulation in emergency department patients with difficult intravenous access. *Ann Emerg Med* 34: 711, 1999.
- Constantino TG, Parikh AK, Satz WA, Fojtik JP: Ultrasonography guided peripheral venous access versus typical approaches in patients with difficult intravenous access. *Ann Emerg Med* 46: 456, 2005.
- Mickiewicz M, Dronen S, Younger J: Central venous catheterization and central venous pressure monitoring in Roberts JR, Hedges JR (eds): *Clinical Procedures in Emergency Medicine*, 4th ed. Philadelphia, Saunders, 2004, p. 413.
- Nomura JT, Leech SJ, Shenbagamurthy S, et al: A randomized controlled trial of ultrasound-assisted lumbar puncture. *J Ultrasound Med* 26: 1341, 2007.
- Ochsner MG, Herr D, Drucker W, et al: A modified Seldinger technique for peritoneal lavage in trauma patients who are obese. *Surg Gynecol Obstet* 173: 158, 1991.
- Uppot RN, Sahani DV, Hahn FP, et al: Impact of obesity on medical imaging and image-guided intervention. *AJR Am J Roentgenol* 188: 433, 2007.

USEFUL WEB RESOURCE

National Center for Health Statistics, Centers for Disease Control and Prevention—<http://www.cdc.gov/nccdphp/dnpa/obesity/trend/index.htm>